

This author's PDF version corresponds to the article as it appeared upon acceptance. Fully formatted PDF versions will be made available soon.

## **Prevalence of disease related malnutrition assessed by NRS2002, STAMP, MNA-SF, one-step and two-step GLIM at admission and discharge: A systematic review and meta-analysis**

doi: 10.6133/apjcn.202605/PP.0003

Published online: May 2026

**Running title:** DRM at admission and discharge

Dongyu Mu MM<sup>1†</sup>, Yi Cheng MM<sup>1†</sup>, Qi Zhang MM<sup>2</sup>, Jie Gong MM<sup>1</sup>, Yana Qi MM<sup>3</sup>, Jiajie Yu PhD<sup>3</sup>, Linan Zeng PhD<sup>4, 5, 6</sup>, Lingli Zhang PhD<sup>3, 4, 5, 6</sup>, Wen Hu MM<sup>1</sup>, Yuan Liu MM<sup>1</sup>, Lei Shi MD<sup>1</sup>

<sup>1</sup>Clinical Nutrition Department, West China Hospital of Sichuan University, Chengdu, Sichuan, China

<sup>2</sup>West China School of Medicine, Sichuan University, Chengdu, Sichuan, China

<sup>3</sup>Chinese Evidence-based Medicine Center, West China Hospital, Sichuan University, Chengdu, Sichuan, China

<sup>4</sup>Department of Pharmacy/Evidence-Based Pharmacy Center, West China Second University Hospital, Sichuan University; Children's Medicine Key Laboratory of Sichuan Province, Chengdu, Sichuan, China

<sup>5</sup>NMPA Key Laboratory for Technical Research on Drug Products In Vitro and In Vivo Correlation, Chengdu, Sichuan, China

<sup>6</sup>Key Laboratory of Birth Defects and Related Diseases of Women and Children, Sichuan University, Ministry of Education, Chengdu, Sichuan, China

<sup>†</sup>Both authors contributed equally to this manuscript

### **Authors' email addresses and contributions:**

DM: dongyumu@scu.edu.cn

Contribution: data collection, data analysis and paper writing.

YC: Miranda1024@wchscu.cn

Contribution: data curation and paper writing.

QZ: zhangqi512@stu.scu.edu.cn

Contribution: data collection and data curation.

JG: gongjie@scu.edu.cn

Contribution: project management.

YQ: qiyana@wchscu.cn

Contribution: methodology.

JY: yujiajie@wchscu.cn

Contribution: methodology.

LNZ: linanzeng@sina.com; ORCID: 0000-0001-9892-2000

Contribution: methodology.

LLZ: zhanglingli@scu.edu.cn

Contribution: supervision.

WH: huw@scu.edu.cn

Contribution: supervision.

YL: liu6341@wchscu.cn

Contribution: conceptualization, funding acquisition and critique providing.

LS: shileilcy@wchscu.cn

Contribution: conceptualization, funding acquisition and critique providing.

**Corresponding Author:** Prof. Yuan Liu, Clinical Nutrition Department, West China Hospital of Sichuan University, 37 Guoxue Alley, Wuhou District, Chengdu 610041, Sichuan Province, China. Tel: +86-189-8060-6459. Email: liu6341@wchscu.cn; Prof. Shi Lei, Clinical Nutrition Department, West China Hospital of Sichuan University, 37 Guoxue Alley, Wuhou District, Chengdu 610041, Sichuan Province, China. Tel: +86-189-8060-3798. Email: shileilcy@wchscu.cn

## ABSTRACT

**Background and Objectives:** Disease-related malnutrition (DRM) mandates routine screening in many nations to ensure quality care as nutritional status often deteriorates during hospitalization. This study aimed to establish tool-specific prevalence benchmarks at admission and discharge, investigate sources of heterogeneity and evaluate diagnostic yields of one-step versus two-step GLIM approaches. **Methods and Study Design:** A systematic search of Embase, PubMed, Web of Science, Cochrane Library, and CINAHL was conducted for cross-sectional studies on nutritional status published between 1 January 2018 and 30 Nov 2025 (PROSPERO ID: CRD42023480467). Eligible studies were pooled in a meta-analysis of proportions using a random-effects model. **Results:** Of 9,693 records retrieved, 179 studies involving 755,092 patients from 30 countries were included. At admission, the pooled prevalence was 0.41 (95% CI 0.35–0.49,  $I^2 = 99.9\%$ ) by NRS2002 ( $\geq 3$ , adults), 0.62 (95% CI 0.57–0.67,  $I^2 = 99.3\%$ ) by MNA-SF ( $\leq 11$ , older adults), 0.55 (95% CI 0.44–0.68,  $I^2 = 98.5\%$ ) by STAMP ( $\geq 2$ , pediatric patients), and 0.39 (95% CI 0.35–0.44,  $I^2 = 99.6\%$ ) by GLIM (adults). At discharge, three studies reported a pooled prevalence of 0.19 (95% CI 0.09–0.40,  $I^2 = 94.6\%$ ) by NRS2002 for adults, significantly lower than admission ( $p = 0.044$ ). Comparing GLIM strategies, no statistically significant difference was observed between one-step and two-step GLIM approaches ( $p = 0.062$ ). **Conclusions:** DRM prevalence remains high at both admission and discharge, varying widely across countries and by tools. Discharge data are critically limited, reflecting gaps in routine screening practices. Regarding diagnosis, findings support a risk-adapted strategy. Pooled estimates may overestimate the true prevalence and should be interpreted with caution.

**Key Words:** nutritional risk, malnutrition, hospitalized patients, prevalence, disease-related malnutrition

## INTRODUCTION

Disease related malnutrition (DRM) occurs when nutrient intake is insufficient or nutrient utilization is impaired during a disease state, leading to weight loss, skeletal muscle loss, physical and mental dysfunctions, and poorer clinical outcomes.<sup>1</sup> This condition is highly prevalent globally and is independently associated with adverse clinical outcomes, including increased infection rates, prolonged hospital stays, and mortality.<sup>2,3</sup> Historically, DRM affects 30–50% of hospitalized patients,<sup>4</sup> with 6%–65% experiencing further deterioration by discharge.<sup>5-9</sup> Given that nutritional support for inpatients who are at nutritional risk or already

malnourished has been shown to improve both nutritional and clinical outcomes, practice guidelines recommend a proactive, screening-based approach to initiating nutritional support during hospitalization.<sup>10</sup>

In pursuit of a world free from all forms of malnutrition, including DRM,<sup>11</sup> WHO established the Sustainable Development Goals (SDGs) and launched the Ambition and Action in Nutrition 2016–2025 initiative.<sup>12, 13</sup> Following these global calls,<sup>14</sup> several countries have implemented national nutrition strategies. For example, China initiated the Healthy China 2030 and National Nutrition Plan (2017–2030),<sup>15, 16</sup> while Lebanon, Vietnam, and Togo published similar plans emphasizing clinical nutrition.<sup>17–20</sup> In parallel, some countries have mandated routine malnutrition screening at hospital admission to trigger subsequent assessment, diagnosis and treatment,<sup>21</sup> aiming to achieve the vision of “hunger-free hospitals”.<sup>22</sup>

In clinical nutrition practice, over 60 screening and diagnostic instruments are currently used, and new tools continue to emerge for specific diseases and settings.<sup>23–26</sup> To ensure comprehensive coverage across all inpatient age groups, we selected four tools that are widely used in routine services and together constitute a representative clinical nutrition workflow: the Nutritional Risk Screening 2002 (NRS2002) and the Global Leadership Initiative on Malnutrition (GLIM) criteria for adults, the Mini Nutritional Assessment–Short Form (MNA-SF) for older adults, and the Screening Tool for the Assessment of Malnutrition in Pediatrics (STAMP) for children.<sup>23–28</sup> These instruments serve distinct roles: NRS2002, MNA-SF, and STAMP are screening tools used to identify patients at nutritional risk who may require further evaluation, while GLIM is a consensus-based diagnostic framework intended to confirm malnutrition based on phenotypic and etiologic criteria.

Despite global and national initiatives, data on DRM prevalence remain fragmented and inconsistent. Reported trends vary widely, with studies showing conflicting trajectories between admission and discharge.<sup>9, 29, 30</sup> Moreover, the comparative impact of one-step (direct diagnosis) versus two-step (screening followed by diagnostic confirmation) GLIM diagnostic approaches on prevalence estimates remains unquantified. A systematic synthesis is therefore essential not only to aggregate these dispersed findings but to establish statistical benchmarks and investigate the sources of variation across diverse global populations.

As many countries approach the midpoint of their national clinical nutrition action plans, tool-specific prevalence estimates offer vital reference points for benchmarking. In China, for instance, screening for nutritional risk within 24 hours of admission is a mandated quality

indicator.<sup>31</sup> Robust, synthesized estimates are therefore needed to establish reliable benchmarks for such routine quality monitoring.

Therefore, we conducted a systematic review and meta-analysis to synthesize the available evidence for benchmarking and to explore global variations. Specifically, this study aimed to: (1) establish tool-specific prevalence benchmarks across countries at admission and discharge; (2) investigate the sources of variation across diverse global populations to assess data consistency; and (3) quantitatively evaluate the diagnostic yields of one-step versus two-step GLIM approaches.

## **MATERIALS AND METHODS**

### ***Study design, protocol registration and reporting guideline compliance***

This was a systematic review and meta-analysis of proportions conducted in accordance with the Reporting Guidelines for Meta-analyses of Observational Studies (MOOSE).<sup>32</sup> This study followed a pre-registered protocol registered in the International Prospective Register of Systematic Reviews (PROSPERO; registration ID: CRD42023480467). To ensure methodological robustness, align with contemporary clinical standards, and address data availability, we implemented specific deviations from the original protocol, including: omitting Subjective Global Assessment (SGA) / Patient-Generated Subjective Global Assessment (PG-SGA) to focus on universal tools (NRS2002, GLIM); updating the timeframe to 2018–2025; reporting prevalence rates exclusively due to data scarcity on screening coverage; and supplementing the Agency for Healthcare Research and Quality (AHRQ) assessment with the Hoy et al. tool.<sup>33</sup> We also restricted inclusion to English-language databases. A detailed summary of all deviations and their scientific justifications is provided in S1 file. As this study involved only secondary analysis of published data, no ethical approval or patient informed consent was required.

### ***Data sources***

We systematically searched five databases for eligible English-language studies published between 1 January 2018 and 30 November 2025, including Embase, PubMed, Web of Science, Cochrane, and CINAHL.

### ***PICOS statement and search strategy***

The brief PICOS statement of the study was as follows: patient problem or population (hospitalized patients), intervention or exposure (nutritional risk or malnutrition with or

without severity grading as defined by the NRS2002, STAMP, MNA-SF; one-step GLIM or two-step GLIM, i.e., any screening tool we selected combined with GLIM at hospital admission or discharge), comparison or control (not applicable), outcome measure (NRS2002, MNA-SF, STAMP, GLIM positive rate at admission or discharge) and study design (cross-sectional studies). For a detailed PICOS overview and search strategies used for each database, along with their original results (Supplementary Table 1 and 2).

### ***Inclusion and exclusion criteria***

We excluded several types of publications from our analysis: editorials, dissertations, books, articles not published in English, conference papers, and news articles. Furthermore, we omitted studies where the full text or nutritional data was inaccessible, even after attempting to contact the authors, as well as those employing alternative designs, and non-original studies, including meta-analyses and reviews. In cases where multiple articles reported on the same population (for instance, when different articles enrolled the same participants but presented results at various time points), we selected only the most recent publication or the one with the largest sample size. Additionally, we excluded non-representative studies, such as those with unclear descriptions of study sites.

### ***Study selection workflow and conflicts settlement***

First, DM conducted a comprehensive database search to export records for each paper, including titles, abstracts, authors, publication years, keywords, etc. Next, the retrieved records were deduplicated using Endnote 21 and Rayyan, then meticulously screened based on predefined protocol. Papers requiring further full-text evaluation were compiled into a single file for subsequent review. Then, two researchers (DM and QZ) independently performed full-text screening, with any discrepancies resolved by senior researchers (YC, YL, and LS).

### ***Data extraction***

The initial draft of the data extraction form was developed by DM. Its final content and format were established after discussions with all researchers and methodologists (YQ, JY, LN, LLZ). Subsequently, two independent investigators (DM and QZ) extracted data from the included studies. Further information was documented in Supplementary Table 3. The primary extracted data encompassed basic article details (e.g., author, publication year, study region, study design, screening/assessment tool, sample size), patient characteristics (e.g.,

mean/median age, female, male), and other relevant data (e.g., timing of nutrition screening/assessment, prevalence of malnutrition at admission/discharge).

### ***Risk of bias assessment***

We assessed the risk of bias and methodological quality of the included studies using two distinct instruments. Study quality was evaluated using AHRQ checklist, with studies categorized as high (8–11 points), fair (4–7 points), or low (0–3 points) quality based on their cumulative scores.

Additionally, we used a validated 10-item tool (Q1-10) specifically designed for prevalence studies by Hoy et al.<sup>33</sup> This tool evaluates four domains: selection, non-response, measurement, and analysis bias. To standardize the application of the 11th item (the overall summary judgment, Q11), we implemented a predefined scoring rule: each preceding item was rated as “yes” (low risk) or “no” (high risk). A study was classified as having a “low risk of bias” if it had fewer than three “no” ratings, “moderate risk” if it had exactly three, and “high risk” if it had four or more. We refined the descriptive criteria for each item to ensure relevance to the specific research context. Two reviewers (DM and QZ) independently performed the assessments immediately following data extraction, with any disagreements resolved through discussion with the broader research team. Detailed assessments are provided in Supplementary Table 4.

### ***Malnutrition screening and diagnostic tools***

The malnutrition screening and diagnostic tools yield different outcome definitions: screening tools (NRS2002, MNA-SF, STAMP) identify “nutritional risk” or “risk of malnutrition,” whereas the diagnostic tool (GLIM) defines “malnutrition.” Accordingly, prevalence estimates were pooled separately within each tool and were not combined across tools, as these outcomes represent distinct conceptual constructs.

#### **NRS2002**

The NRS2002, developed by Kondrup et al. for inpatients aged 19 to 90, was formulated based on 12 randomized controlled trials (RCTs) and retrospectively validated using data from 128 RCTs.<sup>34-36</sup> An NRS2002 score of 3 or higher is called “nutritional risk” and signifies the need for a follow-up nutritional assessment or diagnosis to ascertain whether the patient suffers from malnutrition.

### **MNA-SF**

The MNA-SF, a malnutrition screening tool originally devised for elderly inpatients aged 65 and over,<sup>37,38</sup> has also proven effective in those aged 60 and above.<sup>39</sup> This abbreviated version, consisting of 6 items compared to the full MNA's 30 items,<sup>40</sup> cuts screening time to roughly 5 minutes without sacrificing key predictive factors. Scores of 7 or fewer on the MNA-SF indicate “malnutrition”, whereas scores between 8 and 11 suggest a “risk of malnutrition”. In brief, any score of 11 or less on the MNA-SF necessitates a deeper evaluation.<sup>28</sup>

### **STAMP**

The STAMP serves as a screening tool specifically designed for children and teenagers aged 2 to 17 years in hospital settings.<sup>41</sup> Its initial phase involves a growth assessment, evaluating age-specific height/weight percentiles or Z-scores to pinpoint deviations from expected growth patterns. This aspect differentiates STAMP from other pediatric screening instruments, such as Screening Tool for Risk On Nutritional status and Growth (STRONGkids).<sup>42</sup> A STAMP score of 2 or higher signifies that the patient faces a medium or high “risk of malnutrition”.

### **GLIM**

The GLIM,<sup>27</sup> a diagnostic tool and consensus framework for DRM in adults, emerged in 2019 from the collaborative efforts of five prominent global nutrition societies. Their goal was to standardize the diagnostic criteria for malnutrition.<sup>43</sup> Under GLIM, malnutrition is diagnosed when at least one phenotypic and one etiologic criterion are met. The framework is structured into two main components: an initial screening to pinpoint patients "at risk" through validated tools like NRS2002 or MNA-SF, followed by a diagnostic evaluation to verify malnutrition and determine its severity based on phenotypic criteria. However, in fact, screening may be omitted, leading to the implementation of either the "one-step GLIM" or "two-step GLIM" approach, i.e., one-step GLIM: applying GLIM diagnostic criteria directly to all subjects without prior screening; two-step GLIM: applying GLIM diagnostic criteria only after a positive nutrition risk screening result.<sup>27, 44</sup> For detailed definitions, target populations, and scoring criteria for each of abovementioned four tools, refer to Supplementary Table 5.

### ***Data synthesis and statistical analysis***

We used R (v4.4.2) to perform descriptive analysis and meta-analysis. The prevalence (p') was computed as the number of events (x) divided by the sample size (n) (i.e.,  $p' = x/n$ ). Prior

to pooling, prevalence data underwent log transformation ( $\log p'$ ) to stabilize variance, as the Shapiro–Wilk normality test indicated that log-transformed data exhibited better normality than raw, logit, arcsin, and double arcsin transformations. The pooled prevalence was then derived as an effect size with corresponding 95% confidence intervals (95%CI), using a two-sided significance level of  $\alpha = 0.05$ . A continuity correction was applied by adding 0.5 to  $x$  and 1 to  $n$ , a common practice to handle extreme proportions (e.g., 0% or 100%).<sup>45</sup>

Given the heterogeneity among studies, we adopted a random-effects model utilizing the DerSimonian and Laird approach. Heterogeneity among studies was evaluated using forest plots, Cochran's  $Q$  test, and the  $I^2$  statistic, with  $I^2$  values of 25%, 50%, and 75% indicating low, moderate, and high heterogeneity, respectively ( $p < 0.1$ ). A leave-one-out sensitivity analysis was conducted to assess the stability of the results.

Subgroup analyses and meta-regressions were conducted to explore sources of variability. Potential covariates were pre-specified based on demographic (gender, gender ratio, mean age group), geographic and socioeconomic (region, country, income level), procedural (timing of assessment, assessor type) and methodological quality (individual items from the Hoy et al. and AHRQ tools) factors.

It was followed by univariable meta-regression for each potential covariate in tool groups with more than 10 studies. Covariates that showed statistical significance were then included in the multivariable model. The adjusted pooled prevalences were calculated as model-predicted proportions and converted from the log scale to the original proportion scale, reflecting covariate-adjusted prevalences instead of simple subgroup-specific prevalences.

Publication bias was assessed using funnel plots, Peters' test, and Duval and Tweedie's trim-and-fill method. For outcomes with significant asymmetry ( $p < 0.01$ ), a Bayesian meta-analysis of log-transformed proportions was performed using a normal–normal hierarchical model with a Half-Normal prior on heterogeneity, to obtain robust credible intervals (CrI). This analysis was limited to subgroups with strong evidence of bias and was not part of the primary analysis.

## RESULTS

### *Study selection*

Following Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 (Figure 1), we initially identified 9,693 records. After Endnote 21 and Rayyan removed 2,625 duplicates, 7,068 records were screened by two independent researchers (DM and QZ). After title and abstract screening, the reviewers reached consensus on 729 records, while

6,339 records were excluded as irrelevant. Of the remaining records, 718 full - text reports were assessed for eligibility, with an additional 11 reports being unavailable.

A total of 515 articles were subsequently excluded for the following reasons: 354 were cohort studies, 49 involved irrelevant populations (e.g., outpatients, post-discharge individuals, or community-dwelling populations), 39 focused on tool validation or threshold development, 30 were conference paper, 16 did not provide the necessary numerator and denominator for calculating prevalence, 13 were secondary analysis, 13 were published in Portuguese, Spanish, Russian or French, 11 was a case-control study, 9 only included patients already identified as having nutritional risks, 5 were pilot studies, 5 were interventional and 2 was reviews. After removing 24 repeated reports, 179 studies (represented in 203 reports) were included in the qualitative synthesis, of which 170 were eligible for quantitative analysis.<sup>46-224</sup> Study characteristics were presented in Supplementary Table 6.

### ***Overview of the included studies***

The final analysis included 179 cross-sectional studies published between 2018 and 2025, reporting on 755,092 hospitalized patients across 30 countries. Thirty-one studies were multicenter, and seven reported prevalence at both admission and discharge; only three met both criteria. NRS2002 (n=114) and GLIM (n=82) were the most used tools, followed by MNA-SF (n=36) and STAMP (n=6); sample sizes ranged from 33 to 324,140. Female-to-male ratios were 0.8 (NRS2002), 0.7 (GLIM), 1.0 (MNA-SF), and 0.8 (STAMP). Mean ages were 62.2 years for NRS2002 and 63.1 for GLIM, compared with 74.3 for MNA-SF and 4.7 for STAMP.

By WHO region,<sup>225</sup> 82 studies (45.8%) were from the Western Pacific, 65 (36.3%) from Europe, 15 (8.4%) from the Americas, 12 (6.7%) from the Eastern Mediterranean, and 5 (2.8%) from South-East Asia. According to World Bank classification,<sup>226</sup> 106 studies (59.2%) were conducted in upper middle income countries, 62 (34.6%) in high income countries, and 11 (6.2%) in lower middle income countries.

Across 201 records, 43 department types were represented. The most common were internal medicine and oncology (each 9.0%), followed by general surgery (8.5%), intensive care (7.0%), gastroenterology (6.5%), and geriatrics (5.0%). Grouped by broader categories, these comprised medical specialties (42.3%), surgical specialties (24.9%), oncology (8.5%), critical care (7.5%), geriatrics and palliative care (5.5%), and other specialties (11.4%).

### ***Risk of bias summary***

Methodological quality varied across the 179 included studies. According to the AHRQ checklist, the majority of studies (n = 136, 76.0%) were of fair quality, while 37 (20.7%) were rated as low quality and only six (3.4%) as high quality. When evaluated using the Hoy et al. tool, 73 studies (40.8%) were classified as having a low risk of bias, 55 (30.7%) as moderate risk, and 51 (28.5%) as high risk.

Specific methodological weaknesses were identified in the following areas:

(1) Participant Selection and Representation: 146 studies (81.6%) were at high risk of bias in population representation due to inadequate accounting for demographics such as age and sex. Limitations in sampling frames were noted in 129 studies (72.1%), and 95 (53.1%) relied on convenience sampling rather than consecutive or random selection.

(2) Assessment Timing: 79 studies (44.1%) failed to conduct nutritional screening or assessment within an appropriate timeframe following hospital admission.

(3) Analytical Reporting: Only two studies (1.1%) explicitly described the management of missing data, and six (3.4%) provided adequate details on response rates and data completeness.

(4) Blinding: Reporting of outcome assessor blinding for subjective measurements was rare, occurring in only five studies (2.8%).

### ***Regional and sample size distribution***

Geographic distribution of included studies was uneven. China contributed nearly half the studies for the NRS2002 analysis (k = 42/90, 46.7 %; 47.2% total weight) and was the predominant source for GLIM analysis (k = 34, 42.5 %). However, the random-effects model effectively mitigated the influence of large-sample studies. For adult populations (NRS2002, MNA-SF, GLIM), no single study contributed >5 % to the total weight, indicating pooled estimates were not driven by individual datasets. In the pediatric STAMP analysis, higher individual weights (15.9 % – 17.3 %) reflected the small number of studies (k = 6) rather than sample size disparities. These pooled estimates should be interpreted in the context of this regional dominance.

## ***Prevalence of nutritional risk and malnutrition***

### **Prevalence at admission**

Main results are summarized in Table 1 and detailed in S7 file, with corresponding forest plots shown in Figures 2-5 (detailed forest plots available in Supplementary Figure 1 and 2). Figure 6 illustrates the pooled prevalence at admission across countries. Key findings are highlighted below.

The pooled prevalence of nutritional risk among adult inpatients (NRS2002  $\geq 3$ ) was 0.41 (95% CI, 0.35–0.49), based on 90 studies ( $k = 90$ ). However, substantial heterogeneity was observed ( $I^2 = 99.9\%$ ,  $p < 0.001$ ), indicating significant variation across studies. Similarly, for older adults screened with MNA-SF ( $\leq 11$ ), the pooled prevalence was 0.62 (95% CI 0.57 – 0.67;  $k = 30$ ,  $I^2 = 99.3\%$ ). Pediatric malnutrition risk (STAMP  $\geq 2$ ) was estimated at 0.55 (95% CI 0.44 – 0.68;  $k = 6$ ,  $I^2 = 98.5\%$ ). For adults diagnosed via GLIM criteria, the pooled prevalence was 0.39 (95% CI 0.35 – 0.44;  $k = 80$ ,  $I^2 = 99.6\%$ ).

### ***GLIM subtypes***

One-step GLIM yielded a pooled malnutrition prevalence of 0.42 (95% CI, 0.37 – 0.48;  $k = 44$ ,  $I^2 = 99.1\%$ ). Two-step GLIM showed varying estimates depending on the initial screen: 0.34 (95% CI 0.26 – 0.46;  $k = 18$ ,  $I^2 = 99.7\%$ ) when preceded by NRS2002, and 0.49 (95% CI 0.35 – 0.68;  $k = 7$ ,  $I^2 = 99.3\%$ ) when preceded by MNA-SF. In all cases, the extremely high  $I^2$  values ( $>98\%$ ) suggest that these pooled estimates represent an average of widely dispersed data points rather than a precise global parameter and should be interpreted with caution.

### ***Prevalence at discharge***

Discharge-time data were limited. Pooled NRS2002  $\geq 3$  prevalence at discharge for adult inpatients was 0.19 (95% CI, 0.09 – 0.40;  $k = 3$ ,  $I^2 = 94.6\%$ ), representing a significant decrease from admission ( $p = 0.044$ ). For GLIM-defined malnutrition, only two isolated estimates were available (0.50 for adults via one-step GLIM;<sup>147</sup> 0.33 for older adults via NRS2002-GLIM),<sup>152</sup> precluding meta-analysis. Given the limited number of studies and high heterogeneity, these findings should be viewed as preliminary.

### ***Subgroup analysis and meta-regression***

Subgroup analyses and meta-regressions explored heterogeneity (detailed in Supplementary Table 7 and 8). Given the extreme heterogeneity ( $I^2 >95\%$ ) across most stratifications, reported associations should be interpreted as exploratory trends rather than definitive quantifications.

‘Country’ showed significant subgroup differences ( $p < 0.05$ ) in univariable analysis, though meta-regression did not confirm it as a moderator. In nations with  $>3$  studies, estimates varied widely within instruments (e.g., NRS-2002 prevalence ranged from 0.32 to 0.68 among adults).

Meta-regression identified potential sources of variation, though caution is warranted due to high residual heterogeneity:

NRS2002: A model incorporating sampling frame (Q2) and non-response bias (Q4) explained 54.5% of heterogeneity. High risk of bias in Q4 predicted lower prevalence (ratio 0.40, 95% CI 0.20–0.78;  $p = 0.007$ ), suggesting systematic exclusion of high-risk patients in studies. Conversely, sampling frame deficiencies (Q2) showed a non-significant trend towards overestimation (ratio 1.27;  $p = 0.061$ ).

MNA-SF: A multivariable model explained 78.0% of the heterogeneity (Akaike Information Criterion, AIC = 48.4), identifying age and risk of bias as key factors. Prevalence was higher in patients  $\geq 80$  years compared with those  $\geq 60$  ( $\beta = 0.25$ ,  $p < 0.001$ ). Methodologically, high risk of bias in summary item (Q11) was associated with lower prevalence ( $\beta = -0.30$ ,  $p < 0.001$ ). While this suggests a quantitative reduction, the association primarily indicates bias direction rather than a precise adjustment magnitude.

GLIM & NRS2002–GLIM: High risk of bias in item Q1 (population representation) was consistently associated with higher reported prevalence. For GLIM, Q1 bias explained 49.3% of heterogeneity, associated with a 1.62-fold increase in prevalence ( $\beta = 0.48$ ,  $p < 0.001$ ; 95% CI 1.22–2.14). Similarly, for NRS2002–GLIM, Q1 bias predicted a 1.86-fold increase ( $\beta = 0.62$ ,  $p = 0.049$ ; 95% CI 1.00 – 3.46).

Crucially, despite the explanatory power of these covariates, considerable residual heterogeneity remained, indicating that unmeasured factors—likely specific clinical protocols or local patient case-mix—drive much of the observed variation.

### ***Sensitivity analysis and publication bias***

Leave-one-out sensitivity analyses confirmed that the pooled estimates were robust to individual study exclusion, with no single dataset disproportionately driving the results.

Furthermore, excluding studies from China yielded a pooled NRS2002 prevalence of 44% (95% CI 0.33 – 0.59), slightly higher than the overall estimate (41 %). This indicates that the inclusion of Chinese data did not inflate the global benchmark, but rather contributed to a slightly more conservative estimate.

However, publication bias was detected for NRS2002 (Peters' test  $t = 3.72$ ,  $p < 0.01$ ), GLIM ( $t = 4.38$ ,  $p < 0.01$ ), and one-step GLIM ( $t = 2.39$ ,  $p = 0.02$ ), with visible asymmetry in funnel plots. Trim-and-fill adjustment imputed 43, 34, and 11 missing studies respectively, reducing the pooled prevalence estimates significantly: NRS2002 dropped from 0.41 to 0.22 (46.3% reduction); GLIM from 0.39 to 0.26 (33.3% reduction); and one-step GLIM from 0.42 to 0.35 (16.7% reduction). These substantial adjustments suggest a potential overestimation of prevalence in the published literature due to small-study effects.

## DISCUSSION

### *Prevalence differences of DRM across tools at admission and discharge*

This study synthesized data from 170 studies to benchmark DRM prevalence across varying tools and populations. Based on the available evidence, which is predominantly weighted towards East Asia, our meta-analysis indicates that nutritional risk and malnutrition remain highly prevalent but widely variable.

At admission, 55% of pediatric inpatients were at nutritional risk (STAMP), while in older adults, malnutrition risk reached 62% (MNA-SF). For the general adult population, prevalence estimates were lower, ranging from 39% (GLIM) to 41% (NRS2002). At discharge, data were scarce, with adult nutritional risk estimated at 19% (NRS2002). Crucially, these figures represent an average of highly heterogeneous data rather than uniform global parameters.

Risk was particularly pronounced in older adults. The higher prevalence detected by MNA-SF (62%) versus NRS2002 (53 % in elderly subgroup) aligns with prior comparative research (45% vs. 38%),<sup>227</sup> likely reflecting the MNA-SF's focus on established malnutrition rather than just acute nutritional risk.

While previous reports suggest malnutrition worsens during hospitalization (e.g., 31% vs 36%),<sup>29</sup> our pooled estimates did not reflect this increase. This discrepancy likely stems from data limitations ( $k = 3$  at discharge) and our reliance on cross-sectional comparisons rather than longitudinal monitoring of the same cohort. Regardless of trends, discharge screening remains a critical quality indicator for ensuring effective transitional care. Yet routine

implementation is rare; for instance, Singaporean hospitals lack mandatory discharge screening protocols,<sup>228</sup> potentially explaining the global scarcity of data on this outcome.

### ***Factors and considerations in GLIM-based diagnosis***

When comparing diagnostic strategies, we found no statistical difference between one-step and two-step approaches ( $p = 0.062$ ). This lack of significance is likely attributable to the extreme heterogeneity ( $I^2 >99\%$ ) within subgroups, which widened confidence intervals and masked potential variations.

However, numerical trends indicate a higher yield for the one-step method (42%) compared to the two-step NRS2002–GLIM approach (34%). This mirrors findings from Bian et al.,<sup>44</sup> who attributed the deficit in two-step models to the imperfect sensitivity of screening tools,<sup>229, 230</sup> which create a diagnostic bottleneck. Notably, tool-specific nuances emerged: while NRS2002–GLIM showed a visible numerical reduction compared to one-step (34% vs 42%), MNA-SF–GLIM estimates were nearly identical (39% vs 42%). This suggests that the impact of the bottleneck varies depending on the sensitivity of the initial risk tool.

Crucially, the recent GLIM 5-year update addresses this dichotomy.<sup>27</sup> It explicitly acknowledges that in high-prevalence settings (e.g., oncology, geriatrics), the screening step may be redundant and should be bypassed to avoid missed diagnoses. Conversely, for general hospital populations, the two-step framework remains the pragmatic standard for resource allocation. Given global dietitian shortages (e.g., 1:807 beds in China),<sup>231</sup> using screening to triage low-risk patients is essential for feasibility. Therefore, we advocate the risk-adapted strategy endorsed by the update: one-step diagnosis for high-risk units, and two-step triage for general wards.

### ***Influence of assessor and timing***

Assessor background did not significantly influence pooled estimates overall (S9 file). However, a consistent trend toward higher prevalence was observed when dietitians participated in assessments, particularly for GLIM-based diagnoses. This likely reflects the value of specialized expertise in identifying subtle phenotypic signs (e.g., muscle loss),<sup>232</sup> whereas non-specialists may prioritize acute medical tasks over comprehensive nutritional evaluation, a common barrier driven by time constraints.<sup>233</sup>

Regarding timing, prevalence estimates did not differ significantly across assessment timepoints ( $p > 0.05$ ). While current guidelines generally recommend screening within 24–48 hours of admission,<sup>3, 234, 235</sup> these findings suggest that extending the window to 24–72 hours

may still be acceptable for stable adult inpatients, at least in terms of detection rates, offering valuable flexibility for overburdened wards.

However, earlier assessment remains preferable where feasible. Our data showed that the most substantial numerical prevalence drop (20% relative reduction) occurred when comparing NRS2002 and NRS2002–GLIM within the first 24 hours. This suggests that for this specific pathway, immediate comprehensive assessment is highly effective for rapidly excluding false positives and initiating care.

For critically ill patients, this speed is essential. Nutritional evaluation is ideally completed within 48 hours to allow the timely initiation of enteral nutrition. As guidelines recommend starting feeding within 24–48 hours,<sup>236</sup> aiming to meet approximately 80% of energy and protein requirements within 48–72 hours,<sup>237, 238</sup> delays in identification could compromise these therapeutic targets.

### ***Workforce constraints and future directions***

Yet, achieving such timely compliance is constrained by workforce shortages. Compressing the diagnostic workload into the first 24–48 hours exacerbates the strain on limited staff.<sup>239</sup> Reported staffing ratios highlight this gap: while U.S. guidelines recommend 1 dietitian per 65–75 patients,<sup>240</sup> actual ratios lag significantly, ranging from 1:73–212 in India to approximately 1:807 beds in China (vs recommended 1:150).<sup>231, 241</sup> Even in specialized settings like Canadian NICUs, ratios can vary widely (median 1:25.3).<sup>242</sup>

These constraints underscore the urgent need for automated solutions. Emerging artificial intelligence (AI) tools offer a promising avenue to alleviate manpower limitations.<sup>243</sup> Early applications have already demonstrated feasibility, including the automated calculation of NRS2002 scores via algorithms and machine learning models using electronic health records to predict malnutrition risk.<sup>244, 245</sup> By automating the initial screening or assessment step, AI could enable hospitals to meet tight diagnostic timelines without overwhelming clinical staff or sacrificing accuracy.<sup>246</sup>

Beyond screening, AI shows potential in guiding personalized nutritional therapy. By analyzing inflammatory markers, energy balance, body composition, and disease severity in real time, algorithms can integrate complex clinical signals to predict nutritional trajectories.<sup>247</sup> This capability enables dynamic adjustments to feeding strategies, thereby shortening clinical decision-making cycles and potentially reducing complications such as gastrointestinal intolerance and fluid overload.<sup>248, 249</sup>

### ***Potential overestimation and reference benchmarks***

Given the predominance of data from China, authoritative Chinese multicenter surveys provide a valuable benchmark for assessing bias direction.

For NRS2002, our pooled estimate for China (38%; overall 41%) is similar to the 35.5% reported in the landmark 2008 tertiary hospital survey,<sup>250</sup> but substantially exceeds the 12.0%–17.8% reported in NutritionDay assessments.<sup>127, 251</sup> Crucially, the NutritionDay surveys excluded ICU patients, whereas our meta-analysis incorporated studies with mixed or higher-acuity populations, capturing a sicker segment of the inpatient population compared with general census surveys.

Regarding the overall direction of bias, our analysis presents competing influences but points to a net overestimation. Initial meta-regression suggested that high non-response bias might lead to underestimation due to the exclusion of critically ill patients. However, this effect appears to be outweighed by the inflating factors observed in the benchmark comparison above. Furthermore, publication bias analysis using the trim-and-fill method reduced the estimate to 22%. Yet we must interpret this adjustment with caution: given the extreme heterogeneity ( $I^2 >98\%$ ), the funnel plot asymmetry may reflect small-study effects (e.g., smaller studies focusing on higher-acuity populations) rather than true publication bias, and the trim-and-fill algorithm might impute “spurious” studies and over-correct the pooled estimate. Thus, the adjusted estimate of 22% should be viewed as a conservative lower bound rather than a definitive correction.<sup>252</sup> Therefore, despite the potential for underestimation in specific high-bias studies, the aggregate evidence suggests that the pooled estimate likely overestimates the true prevalence in the general inpatient population.

Similarly, for NRS2002–GLIM, our estimate (34%) is higher than the 12.5% reported in the China Nutrition Fundamental Data 2020 (CNFD 2020) study,<sup>49</sup> which specifically excluded critically ill patients. Therefore, our results likely overrepresent populations with higher clinical acuity rather than the general inpatient census.

For one-step GLIM, the evidence presents a mixed but precautionary picture. The Bayesian meta-analysis remained consistent with the crude estimate (42%), suggesting statistical robustness based on the available data, whereas the trim-and-fill analysis identified potential publication bias and suggested a downward correction of the prevalence to 35%. Given this signal, it is possible that the current pooled estimate for one-step GLIM may overestimate the true prevalence.

### ***Limitations related to high heterogeneity and regional overrepresentation***

A fundamental limitation of this study is the extreme heterogeneity ( $I^2 >95\%$ ) observed across all analyses. Consistent with Cochrane guidance, we explicitly acknowledge that pooling such disparate data carries the risk of producing an invalid average. Consequently, the pooled prevalence rates reported here should not be interpreted as precise global parameters, but strictly as descriptive summaries of the currently available, fragmented literature. In the same vein, the apparent reduction in NRS2002-defined malnutrition from admission to discharge is based on only three highly heterogeneous discharge studies and should therefore be interpreted as a tentative signal rather than firm evidence of a true change over time. By contrast, the comparison between one-step and two-step GLIM strategies did not show a statistically significant difference despite a larger evidence base, and we accordingly treat this as inconclusive rather than proof of equivalence, highlighting the need for further, more standardized data.

Critically, this heterogeneity was not resolved by extensive subgroup stratification or meta-regression. This persistent variability mirrors findings from other meta-analyses,<sup>253, 254</sup> where heterogeneity typically remains extreme despite rigorous adjustment.

However, our analysis enables us to dissect the sources of this variation more granularly. Methodologically, as detailed in our results, study quality indicators—specifically sampling bias (e.g., reliance on single-department convenience sampling) and representativeness bias (e.g., lack of hospital-wide census)—accounted for approximately half of the heterogeneity (54% for NRS2002 and 49% for GLIM). Clinically, variability was driven by four key dimensions (Supplementary Table 8): For NRS2002, (1) hospital setting, with intensive care unit (ICU) prevalence (60%) far exceeding general wards (39%); (2) patient demographics, where older inpatients (age  $\geq 65$ y) show double the risk of younger ones (51% vs 21%); (3) regional disparities, ranging from 39% in Europe to 74% in South-East Asia; for GLIM, (4) economic context, with lower-middle-income nations reporting higher rates (49%) than upper-middle-income ones (35%).

Even after accounting for these factors, substantial unexplained variance remains. This likely stems from the inherent imprecision in defining “malnutrition” itself, coupled with unmeasured variations in local screening protocols driven by administrative mandates, timeframes, and economic levers. For instance, China and Portugal not only enforce mandatory screening via national health commissions as a quality indicator, but also stipulate distinct time windows (e.g., within 24 hours in China versus 48 hours in Portugal), effectively broadening the screened denominator and capturing different patient profiles.<sup>14, 255-257</sup>

Furthermore, reimbursement policies in China and Switzerland link insurance payments directly to positive screening results (NRS2002  $\geq 3$ ), creating strong economic incentives for rigorous identification.<sup>257, 258</sup> Thus, we interpret these variations as a reflection of the real-world diversity in clinical nutrition practice.

Furthermore, studies from China contributed nearly half of the data for NRS2002 and GLIM analyses. While our random-effects model mitigated the statistical dominance of large-sample studies (no single study weighted  $>5\%$ ), the pooled estimates inevitably reflect the epidemiological characteristics of East Asian inpatient populations more heavily. Data from the African Region remain scarce, limiting the generalizability of our benchmarks to these settings.

### ***Conclusions***

The prevalence of DRM among hospitalized patients remains high at both admission and discharge, though estimates vary across countries and by the screening and diagnostic tool used. Discharge-time data were rarely reported, likely reflecting limited implementation of routine discharge screening in clinical practice.

Regarding diagnosis, GLIM-based diagnostic approaches showed no significant differences between one-step and two-step strategies. Consequently, our findings support the adoption of a risk-adapted strategy: one-step diagnosis for high-risk units to ensure detection, and two-step triage for general wards to optimize limited workforce resources.

Methodologically, these meta-analytic estimates may tend to overestimate the true prevalence and should be extrapolated with caution. Further observational studies assessing nutritional status at both admission and discharge are needed to better capture in-hospital nutritional trends and support nutrition-related quality assurance efforts.

### **SUPPLEMENTARY MATERIALS**

All supplementary tables and figures are available upon request from the editorial office, and are also accessible on the journal's webpage ([apjcn.qdu.edu.cn](http://apjcn.qdu.edu.cn)).

### **ACKNOWLEDGEMENTS**

We thank the authors of the primary studies included in this review for their valuable contributions to the field.

## CONFLICT OF INTEREST AND FUNDING DISCLOSURE

The authors declare no conflict of interest.

This research was funded by the Key Research and Development Project of the Ministry of Science and Technology Project (No.2023YFF1104405).

## REFERENCES

1. Cederholm T, Bosaeus I. Malnutrition in Adults. *N Engl J Med*. 2024;391:155–65. doi: 10.1056/NEJMra2212159.
2. ESPEN. ESPEN Fact Sheet of Disease Related Malnutrition (DRM). 2017/2/1 [cited 2025/8/7]; Available from: <https://www.espen.org/images/files/ESPEN-Fact-Sheets/ESPEN-Fact-Sheet-Disease-Related-Malnutrition-DRM.pdf>
3. Seron-Arbeloa C, Labarta-Monzon L, Puzo-Foncillas J et al. Malnutrition Screening and Assessment. *Nutrients*. 2022;14(12):2392. doi: 10.3390/nu14122392.
4. WHO Regional Office for Europe. Disease-related malnutrition: a time for action. 2023/1/1 [cited 2025/8/8]; Available from: <https://iris.who.int/handle/10665/375033>
5. McWhirter JP, Pennington CR. Incidence and recognition of malnutrition in hospital. *BMJ*. 1994;308:945–48. doi: 10.1136/bmj.308.6934.945.
6. Lean M, Wiseman M. Malnutrition in hospitals. *BMJ*. 2008;336:290. doi: 10.1136/bmj.39449.723090.80.
7. Kondrup J, Allison SP, Elia M, Vellas B, Plauth M. ESPEN guidelines for nutrition screening 2002. *Clin Nutr*. 2003;22:415–21. doi: 10.1016/s0261-5614(03)00098-0.
8. Zhu M, Wei J, Chen W, Yang X, Cui H, Zhu S. Nutritional Risk and Nutritional Status at Admission and Discharge among Chinese Hospitalized Patients: A Prospective, Nationwide, Multicenter Study. *J Am Coll Nutr*. 2017;36:357–63. doi: 10.1080/07315724.2017.1304293.
9. Cass AR, Charlton KE. Prevalence of hospital-acquired malnutrition and modifiable determinants of nutritional deterioration during inpatient admissions: A systematic review of the evidence. *J Hum Nutr Diet*. 2022;35:1043–58. doi: 10.1111/jhn.13009.
10. Gomes F, Baumgartner A, Bounoure L et al. Association of Nutritional Support With Clinical Outcomes Among Medical Inpatients Who Are Malnourished or at Nutritional Risk: An Updated Systematic Review and Meta-analysis. *JAMA Netw Open*. 2019;2(11):e1915138. doi: 10.1001/jamanetworkopen.2019.15138.
11. Barker LA, Gout BS, Crowe TC. Hospital malnutrition: prevalence, identification and impact on patients and the healthcare system. *Int J Environ Res Public Health*. 2011;8:514–27. doi: 10.3390/ijerph8020514.
12. WHO. Ambition and Action in Nutrition 2016-2025. 2017/5/18 [cited 2025/2/22]; Available from: <https://www.who.int/publications/i/item/9789241512435>
13. WHO. SDG Target 2.2 Malnutrition. 2025/1/30 [cited 2025/2/22]; Available from: [https://www.who.int/data/gho/data/themes/topics/sdg-target-2\\_2-malnutrition](https://www.who.int/data/gho/data/themes/topics/sdg-target-2_2-malnutrition)

14. Cai L, Hu X, Liu S et al. China is implementing the national nutrition plan of action. *Front Nutr.* 2022;9:983484. doi: 10.3389/fnut.2022.983484.
15. General Office of the State Council of the People's Republic of China. National Nutrition Plan (2017-2030). 2017/7/13 [cited 2026/2/3]; Available from: [https://www.gov.cn/zhengce/content/2017-07/13/content\\_5210134.htm](https://www.gov.cn/zhengce/content/2017-07/13/content_5210134.htm)
16. General Office of the State Council of the People's Republic of China. Healthy China 2030. 2016/10/25 [cited 2025/2/22]; Available from: [https://www.gov.cn/zhengce/2016-10/25/content\\_5124174.htm](https://www.gov.cn/zhengce/2016-10/25/content_5124174.htm)
17. Lebanon. National Nutrition Strategy and Action Plan 2021 - 2026. 2021/6/1 [cited 2025/2/24]; Available from: <https://www.fao.org/faolex/results/details/en/c/LEX-FAOC216967/>
18. Togo. Plan Stratégique National multisectoriel de la nutrition au Togo (2019-2023). 2019/1/1 [cited 2025/8/4]; Available from: <https://www.fao.org/faolex/results/details/en/c/LEX-FAOC223827/>
19. Viet Nam. Decision 718/QD-BYT 2018 approving the National Action Plan on Nutrition by 2020. 2018/1/29 [cited 2025/2/24]; Available from: <https://www.fao.org/faolex/results/details/en/c/LEX-FAOC230565/>
20. Viet Nam. Decision 1294/QD-BYT 2022 issuing the Action Plan to implement the National Nutrition Strategy to 2025. 2022/4/19 [cited 2025/2/24]; Available from: <https://www.fao.org/faolex/results/details/en/c/LEX-FAOC230025/>
21. de van der Schueren MAE, Jager-Wittenaar H. Malnutrition risk screening: New insights in a new era. *Clin Nutr.* 2022;41:2163–68. doi: 10.1016/j.clnu.2022.08.007.
22. Xin W, Yang Y, Shi H-P. Hunger-free hospital initiative: fighting “the skeleton in the hospital closet”. *Precis Nutr.* 2022;1(3):e00018. doi: 10.1097/pn9.000000000000018.
23. Cortes-Aguilar R, Malih N, Abbate M, Fresneda S, Yanez A, Bennasar-Veny M. Validity of nutrition screening tools for risk of malnutrition among hospitalized adult patients: A systematic review and meta-analysis. *Clin Nutr.* 2024;43:1094–116. doi: 10.1016/j.clnu.2024.03.008.
24. Oishi K, Inoue-Hamano A, Hamada Y. Development of a New Malnutrition Screening Tool for Patients: Human Key Tool of Nutrition. *Am J Med Open.* 2025;13:100086. doi: 10.1016/j.ajmo.2024.100086.
25. Shi J, Liu T, Ge Y et al. Cholesterol-modified prognostic nutritional index (CPNI) as an effective tool for assessing the nutrition status and predicting survival in patients with breast cancer. *BMC Med.* 2023;21(1):512. doi: 10.1186/s12916-023-03225-7.
26. Foo J, Roberts M, Williams LT, Osadnik C, Bauer J, O'Shea MC. An Automated Malnutrition Screening Tool Using Routinely Collected Data for Older Adults in Long-Term Care: Development and Internal Validation of AutoMal. *J Am Med Dir Assoc.* 2026;27(2):105252. doi: 10.1016/j.jamda.2024.105252.
27. Cederholm T, Jensen GL, Correia M et al. The GLIM consensus approach to diagnosis of malnutrition: A 5-year update. *Clin Nutr.* 2025;49:11–20. doi: 10.1016/j.clnu.2025.03.018.
28. Reber E, Gomes F, Vasiloglou MF, Schuetz P, Stanga Z. Nutritional Risk Screening and Assessment. *J Clin Med.* 2019;8(7):1065. doi: 10.3390/jcm8071065.

29. van Vliet IMY, Gomes-Neto AW, de Jong MFC, Jager-Wittenaar H, Navis GJ. High prevalence of malnutrition both on hospital admission and predischARGE. *Nutrition*. 2020;77:110814. doi: 10.1016/j.nut.2020.110814.
30. Poudineh S, Shayesteh F, Kermanchi J et al. A multi-centre survey on hospital malnutrition: result of PNSI study. *Nutr J*. 2021;20(1):87. doi: 10.1186/s12937-021-00741-1.
31. National Clinical Nutrition Quality Control Center. National Report on the Services, Quality and Safety in Medical Care System (2020): Clinical Nutrition. Tianjin: People's Medical Publishing House; 2023.
32. Brooke BS, Schwartz TA, Pawlik TM. MOOSE Reporting Guidelines for Meta-analyses of Observational Studies. *JAMA Surg*. 2021;156:787–88. doi: 10.1001/jamasurg.2021.0522.
33. Hoy D, Brooks P, Woolf A et al. Assessing risk of bias in prevalence studies: modification of an existing tool and evidence of interrater agreement. *J Clin Epidemiol*. 2012;65:934–39. doi: 10.1016/j.jclinepi.2011.11.014.
34. Cederholm T, Barazzoni R, Austin P et al. ESPEN guidelines on definitions and terminology of clinical nutrition. *Clin Nutr*. 2017;36:49–64. doi: 10.1016/j.clnu.2016.09.004.
35. Kondrup J, Rasmussen HH, Hamberg O, Stanga Z, Ad Hoc EWG. Nutritional risk screening (NRS 2002): a new method based on an analysis of controlled clinical trials. *Clin Nutr*. 2003;22:321–36. doi: 10.1016/s0261-5614(02)00214-5.
36. Kang Y. Nutritional risk screening and malnutrition diagnosis. *China Medical News*. 2021;36(11):18. doi: 10.3760/cma.j.issn.1000-8039.2021.11.126.
37. Volkert D, Beck AM, Cederholm T et al. ESPEN guideline on clinical nutrition and hydration in geriatrics. *Clin Nutr*. 2019;38:10–47. doi: 10.1016/j.clnu.2018.05.024.
38. Rubenstein LZ, Harker JO, Salva A, Guigoz Y, Vellas B. Screening for undernutrition in geriatric practice: developing the short-form mini-nutritional assessment (MNA-SF). *J Gerontol A Biol Sci Med Sci*. 2001;56(6):M366–72. doi: 10.1093/gerona/56.6.m366.
39. Kaiser MJ, Bauer JM, Ramsch C et al. Validation of the Mini Nutritional Assessment short-form (MNA-SF): a practical tool for identification of nutritional status. *J Nutr Health Aging*. 2009;13:782–88. doi: 10.1007/s12603-009-0214-7.
40. Guigoz Y, Vellas B, Garry PJ. Assessing the nutritional status of the elderly: The Mini Nutritional Assessment as part of the geriatric evaluation. *Nutr Rev*. 1996;54(1 Pt 2):S59–65. doi: 10.1111/j.1753-4887.1996.tb03793.x.
41. McCarthy H, McNulty H, Dixon M, Eaton - Evans MJ. Screening for nutrition risk in children: the validation of a new tool. *J Hum Nutr Diet*. 2008;21:395 – 96. doi: 10.1111/j.1365-277X.2008.00881\_31.x.
42. Hulst JM, Zwart H, Hop WC, Joosten KF. Dutch national survey to test the STRONGkids nutritional risk screening tool in hospitalized children. *Clin Nutr*. 2010;29:106–11. doi: 10.1016/j.clnu.2009.07.006.

43. Jensen GL, Cederholm T, Correia M et al. GLIM Criteria for the Diagnosis of Malnutrition: A Consensus Report From the Global Clinical Nutrition Community. *JPEN J Parenter Enteral Nutr.* 2019;43:32–40. doi: 10.1002/jpen.1440.
44. Bian W, Li Y, Wang Y et al. Prevalence of malnutrition based on global leadership initiative in malnutrition criteria for completeness of diagnosis and future risk of malnutrition based on current malnutrition diagnosis: systematic review and meta-analysis. *Front Nutr.* 2023;10:1174945. doi: 10.3389/fnut.2023.1174945.
45. Sutton AJ, Abrams KR, Jones DR, Sheldon TA, Song F. *Methods for meta-analysis in medical research.* Chichester: Wiley; 2000.
46. Rajan S, Dhamija E, Malhotra N et al. Nutritional assessment in gynecological cancers: Experience from an Indian tertiary care center. *Eur J Obstet Gynecol Reprod Biol.* 2025;310:113964. doi: 10.1016/j.ejogrb.2025.113964.
47. Ozsenel EB, Kahveci G, Pekcioglu Y, Guner B, Basat S. The Overlooked Threat of Malnutrition: A Point Prevalence Study Based on NRS-2002 Screening in a Tertiary Care Hospital. *J Clin Med.* 2025;14(11):3976. doi: 10.3390/jcm14113976.
48. Munk T, Holmsted SF, Beck AM, Laursen L, Rasmussen HH, Knudsen AW. The effect of a digital energy- and protein-dense food concept with a nutritional tracker on dietary intake in hospitalized patients at nutritional risk - A one-day cross-sectional study. *Clin Nutr ESPEN.* 2025;68:523–29. doi: 10.1016/j.clnesp.2025.05.037.
49. Liu Y, Song C, Wang X et al. Prevalence of malnutrition among adult inpatients in China: a nationwide cross-sectional study. *Sci China Life Sci.* 2025;68:1487–97. doi: 10.1007/s11427-023-2619-7.
50. Liu JF, Jiang QX, Liu J, Liu AL, Wang YH. Investigation and Analysis of Frailty and Nutritional Status in Patients With Inflammatory Bowel Disease. *Crohns Colitis 360.* 2025;7(2):otaf010. doi: 10.1093/crocol/otaf010.
51. Lehmann U, Uhlmann K, Meichtry A et al. Malnutrition and sarcopenia in inpatient rehabilitation: prevalence and associations with changes in bodyweight, muscle strength, and functional independence. *J Rehabil Med.* 2025;57:jrm42215. doi: 10.2340/jrm.v57.42215.
52. Kong X, Liu P, Wang G, Sun S, Li L. Methods for diagnosing malnutrition in patients with esophageal cancer, and the association with nutritional and inflammatory indices: A cross-sectional study. *Oncol Lett.* 2025;29(5):223. doi: 10.3892/ol.2025.14969.
53. Mphwanthe G, Columbus T, Wasambo L et al. Malnutrition prevalence at hospital admission and discharge based on the two-step Global Leadership Initiative on Malnutrition: A prospective observational study in Malawian adult patients. *Nutr Clin Pract.* 2025. doi: 10.1002/ncp.11347.
54. Doganay M, Halil MG, Uyar M et al. Prevalence of malnutrition risk in hospitalized patients: a large nationwide study. *J Health Popul Nutr.* 2025;44(1):137. doi: 10.1186/s41043-025-00891-6.

55. Alamri A, Alaamer K, Almogbel Y et al. Prevalence of Malnutrition in People Hospitalized for Surgery: Prospective Cross-Sectional Study. *Healthcare*. 2025;13(4):380. doi: 10.3390/healthcare13040380.
56. Zhou L, Ding Z, Wang Q, Wu R, Jin K. Evaluation of malnutrition by objective nutritional indexes and predictors in hospitalized patients with COVID-19. *J Clin Biochem Nutr*. 2024;75:153–60. doi: 10.3164/jcbtn.24-73.
57. Zhou B, Zhang Y, Hiesmayr M et al. Dietary Provision, GLIM-Defined Malnutrition and Their Association with Clinical Outcome: Results from the First Decade of nutritionDay in China. *Nutrients*. 2024;16(4):569. doi: 10.3390/nu16040569.
58. Yogesh M, Patel J, Makwana N, Mody M. The triad of physiological challenges: investigating the intersection of sarcopenia, malnutrition, and malnutrition-sarcopenia syndrome in patients with COPD - a cross-sectional study. *BMC Pulm Med*. 2024;24(1):71. doi: 10.1186/s12890-024-02884-3.
59. Tu Y, Chen F, Yu Q, Song L, Chen M. Application of NRS2002 and PG-SGA in nutritional assessment for perioperative patients with head and neck squamous cell carcinoma: An observational study. *Medicine*. 2024;103(43):e40025. doi: 10.1097/MD.00000000000040025.
60. Wang R, Huang L, Xu M, Yu X, Wang H. Comparison of different nutritional screening tools in nutritional screening of patients with cirrhosis: A cross-sectional observational study. *Heliyon*. 2024;10(10):e30339. doi: 10.1016/j.heliyon.2024.e30339.
61. Slettahjell HB, Bastakis M, Biering-Sorensen F, Strom V, Henriksen C. Defining malnutrition in persons with spinal cord injury - does the Global Criteria for Malnutrition work? *Food Nutr Res*. 2024;68:9989. doi: 10.29219/fnr.v68.9989.
62. Tan S, Jiang J, Qiu L et al. Prevalence of Malnutrition in Patients with Hepatocellular Carcinoma: A Comparative Study of GLIM Criteria, NRS2002, and PG-SGA, and Identification of Independent Risk Factors. *Nutr Cancer*. 2024;76:335–44. doi: 10.1080/01635581.2024.2314317.
63. Ruiz-Rosso R, Moreno-Camara S, Gutierrez-Sanchez B, da-Silva-Domingues H, Del-Pino-Casado R, Palomino-Moral PA. Factors Influencing Nutritional Status in Hospitalized Individuals Aged 70 and Above. *Nutrients*. 2024;16(5):645. doi: 10.3390/nu16050645.
64. Omar OM, Tayel DI, Saleh RM, Ramadan MA. The malnutrition screening tool STRONGKIDS performed better than other screening tools for children hospitalised in Egypt. *Acta Paediatr*. 2024;113:1006–12. doi: 10.1111/apa.17123.
65. Oner O, Hanci V, Gurkok MC et al. The impact of malnutrition and frailty on morbidity and mortality in geriatric internal intensive care patients: a prospective study. *Turk Geriatr Derg*. 2024;27:135–45. doi: 10.29400/tjgeri.2024.387.
66. Ma W, Cai B, Li HX et al. GLIM-defined malnutrition in patients with acute abdomen associated with poor prognosis and increased economic burden: A cross-sectional study. *Nutr Clin Pract*. 2024;39:1364–74. doi: 10.1002/ncp.11202.
67. Mercan Baspinar M, Guden A, Calik M, Mavis O, Kamat S, Basar H. Determination of malnutrition, frailty, polypharmacy, multimorbidity, and mortality risk in hospitalized elderly COVID-19 pneumonia

- cases with and without possible sarcopenia. *Turk Geriatr Derg.* 2024;27:239–47. doi: 10.29400/tjgeri.2024.397.
68. Reyes L M, Garcia-Moreno R M, Lopez-Plaza B, Milla S P. Prevalence, Risk Factors, and Clinical Management of Disease-Related Malnutrition in Hospitalized Patients: A Descriptive Analysis Using GLIM and SGA Criteria. *Nutrients.* 2024;16(23):4099. doi: 10.3390/nu16234099.
69. Jazinaki MS, Norouzy A, Arabi SM, Moghadam M, Esfahani AJ, Safarian M. Two-step GLIM approach using NRS-2002 screening tool vs direct GLIM criteria application in hospital malnutrition diagnosis: A cross-sectional study. *Nutr Clin Pract.* 2024;39:1419–30. doi: 10.1002/ncp.11229.
70. Yang JY, Zhu JY, Cao PL et al. Applicability of Global Leadership Initiative on Malnutrition (GLIM) criteria in nutrition risk screening for patients over 60 years old with digestive system tumors - A retrospective study. *Nutr Clin Metab.* 2024;38:194–202. doi: 10.1016/j.nupar.2024.06.002.
71. Zuo J, Huang Y, Huang Z et al. Comparison of three objective nutritional screening tools for identifying GLIM-defined malnutrition in patients with gastric cancer. *Eur J Clin Nutr.* 2025;79:64–70. doi: 10.1038/s41430-024-01514-9.
72. Wang J, Xu Q, Xie H et al. Comparison of the Global Leadership Initiative on Malnutrition and the Patient-Generated Subjective Global Assessment for diagnosing malnutrition in patients undergoing surgery for hepatobiliary and pancreatic malignancies. *Nutr Hosp.* 2024;41:835–42. doi: 10.20960/nh.05056.
73. Ottestad I, Thoresen L, Tangvik RJ et al. Inadequate nutritional care for malnourished patients in four university hospitals - The QuaNuT study. *Clin Nutr Open Sci.* 2024;54:63–77. doi: 10.1016/j.nutos.2024.01.002.
74. Qing H, Zhang X D, Yang E et al. Nutritional status and nutritional intervention of older inpatients in China. *J Nutr Health Aging.* 2024;28(3):100169. doi: 10.1016/j.jnha.2024.100169.
75. Godala M, Gaszynska E, Walczak K, Malecka-Wojcieszko E. An Evaluation of the Usefulness of Selected Screening Methods in Assessing the Risk of Malnutrition in Patients with Inflammatory Bowel Disease. *Nutrients.* 2024;16(6):814. doi: 10.3390/nu16060814.
76. Marquez Mesa E, Guerra Cabrera AJ, Gomez de Segura IL, Suarez Llanos JP. Comparison of CIPA Nutritional Screening with GLIM Criteria for Malnutrition, Prognostic Evolution, and Association with Phase Angle in Hospitalized Patients. *Nutrients.* 2024;16(21):3652. doi: 10.3390/nu16213652.
77. Cortes R, Yanez AM, Capitan-Moyano L, Millan-Pons A, Bennasar-Veny M. Evaluation of different screening tools for detection of malnutrition in hospitalised patients. *J Clin Nurs.* 2024;33:4759–71. doi: 10.1111/jocn.17170.
78. Clement D, van Leerdam ME, Tesselaar MET et al. The global leadership into malnutrition criteria reveals a high percentage of malnutrition which influences overall survival in patients with gastroenteropancreatic neuroendocrine tumours. *J Neuroendocrinol.* 2024;36(4):e13376. doi: 10.1111/jne.13376.

79. Guan C, Lu T, Liao Z et al. Nutritional Status and Incidence of Radiation-Induced Oral Mucositis in Nasopharyngeal Carcinoma Patients Treated with Chemoradiotherapy. *Nutr Cancer*. 2024;76:196–206. doi: 10.1080/01635581.2023.2294523.
80. Alvarez-Altamirano K, Bejarano-Rosales MP, Gonzalez-Rodriguez BK et al. Prevalence of nutritional risk and malnutrition in hospitalized patients: a retrospective, cross-sectional study of single-day screening. *Appl Physiol Nutr Metab*. 2024;49:838–43. doi: 10.1139/apnm-2023-0190.
81. Pham Thi Lan A, Le Ngoc Thao V, Doan Duy T, Pham Nhat T. Malnutrition and other associated factors among the hospitalized elderly in Vietnam: A cross-sectional study. *Medicine*. 2024;103(36):e39563. doi: 10.1097/MD.00000000000039563.
82. Zhou LM, Fu JY, Ding Z, Jin KM, Wu RJX, Ye LX. Comparison of GLIM, SGA, PG-SGA, and PNI in diagnosing malnutrition among hepatobiliary-pancreatic surgery patients. *Front Nutr*. 2023;10:1116243. doi: 10.3389/fnut.2023.1116243.
83. Zheng J, Yu FM, Chen YY, Ming F, Rao ZY. The Practice of Nutrition Risk Screening, Nutrition Status Assessment, and Nutrition Support Action (NRASA) in Hospitalized Patients. *Top Clin Nutr*. 2023;38:249–61. doi: 10.1097/tin.0000000000000311.
84. Yang W, Guo G, Mao L et al. Comparison of the GLIM criteria with specific screening tool for diagnosing malnutrition in hospitalized patients with cirrhosis: A descriptive cross-sectional study. *JPEN J Parenter Enteral Nutr*. 2023;47:310–21. doi: 10.1002/jpen.2452.
85. Yang W, Guo G, Cui B et al. Malnutrition according to the Global Leadership Initiative on Malnutrition criteria is associated with in-hospital mortality and prolonged length of stay in patients with cirrhosis. *Nutrition*. 2023;105:111860. doi: 10.1016/j.nut.2022.111860.
86. Wei W, Yan PG, Wang F et al. Malnutrition Defined by the Global Leadership Initiative on Malnutrition (GLIM) Criteria in Hospitalized Patients with Ulcerative Colitis and Its Association with Clinical Outcomes. *Nutrients*. 2023;15(16):3572. doi: 10.3390/nu15163572.
87. Wang W, Dai L, Ma J, Gu L, Xie H, Fu J. Malnutrition accelerates the occurrence of infectious complications in patients with chronic kidney disease: A cross-sectional survey of 682 patients with chronic kidney disease. *Nutr Clin Pract*. 2023;38:1167–74. doi: 10.1002/ncp.11040.
88. Wang L, Li P, Hu Y et al. Relationship between preoperative malnutrition, frailty, sarcopenia, body composition, and anthropometry in elderly patients undergoing major pancreatic and biliary surgery. *Front Nutr*. 2023;10:1135854. doi: 10.3389/fnut.2023.1135854.
89. Wang JJ, Zhuang QL, Tan SJ et al. Loss of body weight and skeletal muscle negatively affect postoperative outcomes after major abdominal surgery in geriatric patients with cancer. *Nutrition*. 2023;106:111907. doi: 10.1016/j.nut.2022.111907.
90. Thomsen TK, Pedersen JL, Sloth B, Damsgaard EM, Rud CL, Hvas CL. Nutritional risk screening in a Danish university hospital is insufficient and may underestimate nutritional risk: A cross-sectional study. *J Hum Nutr Diet*. 2023;36:108–15. doi: 10.1111/jhn.13025.

91. Silva RJG, Grippa WR, Neto L et al. Factors Associated with the Nutritional Status of Women with Non-Metastatic Breast Cancer in a Brazilian High Complexity Oncology Center. *Nutrients*. 2023;15(23):4961. doi: 10.3390/nu15234961.
92. Poveda MD, Suárez-de-la-Rica A, Minchot EC et al. The Prevalence and Impact of Nutritional Risk and Malnutrition in Gastrointestinal Surgical Oncology Patients: A Prospective, Observational, Multicenter, and Exploratory Study. *Nutrients*. 2023;15(14):3283. doi: 10.3390/nu15143283.
93. Ozorio GA, Ribeiro LMK, Santos BC et al. Exploring the use of the GLIM criteria to diagnose malnutrition in cancer inpatients. *Nutrition*. 2023;116:112195. doi: 10.1016/j.nut.2023.112195.
94. Ouaijan K, Hwalla N, Kandala NB, Mpinga EK. Prevalence of Malnutrition in Hospitalized Patients in Lebanon Using Nutrition Risk Screening (NRS-2002) and Global Leadership Initiative on Malnutrition (GLIM) Criteria and Its Association with Length of Stay. *Healthcare*. 2023;11(5):730. doi: 10.3390/healthcare11050730.
95. Orell H, Pohju A, Tuokkola J et al. Time to act!-A cross-sectional study on how nutritional risk increases during hospitalization and associates with worse outcome. *Clin Nutr ESPEN*. 2023;57:364–74. doi: 10.1016/j.clnesp.2023.07.016.
96. Nucci D, Gianfredi V, Ferrara P et al. Association between Malnutrition and Depression in Patients with Cancer: The Importance of Nutritional Status Evaluation in Cancer Care. *Int J Environ Res Public Health*. 2023;20(3):2295. doi: 10.3390/ijerph20032295.
97. Nozawa Y, Nishitani-Yokoyama M, Shimada K et al. Nutritional Status in Patients Undergoing Phase II Cardiac Rehabilitation by Mini Nutritional Assessment. *Cardiol Res*. 2023;14:133–41. doi: 10.14740/cr1479.
98. Nguyen LHT, Dang AK, Tran TV et al. The role of nutritional risk evaluation in predicting adverse outcomes among patients with severe COVID-19 in Vietnam. *Front Nutr*. 2023;10:1245816. doi: 10.3389/fnut.2023.1245816.
99. Naaman RK, Bakhsh MA, Alsherebi RK, Sultan RM. Prevalence and Recognition of Malnutrition in Elderly Patients Admitted to a University Hospital in Jeddah, Saudi Arabia. *Top Clin Nutr*. 2023;38:318–26. doi: 10.1097/tin.0000000000000344.
100. Meissner C, Tiegges S, Broehl M, Otto R, Ridwelski K. International study on the prevalence of malnutrition in centralized care for colorectal cancer patients. *Innov Surg Sci*. 2023;8(2):83–92. doi: 10.1515/iss-2023-0015.
101. Li Y, Liu F, Xie H, Zhu Y. Investigation and analysis of frailty and nutrition status in older adult patients with hip fracture. *Nutr Clin Pract*. 2023;38:1063–72. doi: 10.1002/ncp.10993.
102. Leandro-Merhi VA, Dos Santos HAV, Almendra AAR, de Aquino JLB. Nutritional indicators' performance in malnutrition diagnosis of hospitalized elderly patients. *Exp Gerontol*. 2023;181:112286. doi: 10.1016/j.exger.2023.112286.
103. Kaya DG, Akin ZC, Orucoglu B, Celik E. Comparison of four different nutritional risk screening tools in hospitalized children. *Rev Nutr*. 2023;36:e220239. doi: 10.1590/1678-9865202235e220239.

104. Kang LH, Liu XL, Ji W et al. Association of Neutrophil-to-Lymphocyte Ratio with Nutrition in Patients with Various Types of Malignant Tumors: A Multicenter Cross-Sectional Study. *J Inflamm Res.* 2023;16:1419–29. doi: 10.2147/jir.S401189.
105. Feng G, Liu M, Li CY et al. The influence of the China GLIM standards on the diagnosis of malnutrition in patients with hematopoietic stem cell transplant. *Front Nutr.* 2022;9:1077442. doi: 10.3389/fnut.2022.1077442.
106. Elif Öztürk M, Yabancı Ayhan N. Evaluation of malnutrition and screening tools in hospitalized children. *Clin Nutr ESPEN.* 2023;57:770–78. doi: 10.1016/j.clnesp.2023.08.031.
107. Deng LH, Chi K, Zong Y, Li Y, Chen MG, Chen P. Malnutrition in patients with head and neck cancer undergoing radiotherapy: A cross-sectional study. *Eur J Oncol Nurs.* 2023;66:102387. doi: 10.1016/j.ejon.2023.102387.
108. Chen X, Liu X, Ji W et al. The PG-SGA outperforms the NRS 2002 for nutritional risk screening in cancer patients: a retrospective study from China. *Front Nutr.* 2023;10:1272420. doi: 10.3389/fnut.2023.1272420.
109. Chen W, Ding Q, Zhang SK et al. Nutritional status in patients with active pulmonary tuberculosis and new nutritional risk screening model for active tuberculosis: a national, multicenter, cross-sectional study in China. *J Thorac Dis.* 2023;15:2779–99. doi: 10.21037/jtd-23-623.
110. Cervera-Díaz MDC, López-Gómez JJ, García-Virto V, Aguado-Hernández HJ, De Luis-Román DA. Prevalence of sarcopenia in patients older than 75 years admitted for hip fracture. *Endocrinol Diabetes Nutr (Engl Ed).* 2023;70:396–407. doi: 10.1016/j.endien.2023.06.001.
111. Carretero Gómez J, Galeano Fernández TF, Vidal Ríos AS et al. Malnutrition and sarcopenia worsen short- and long-term outcomes in internal medicine inpatients. *Postgrad Med J.* 2023;99(1168):56–62. doi: 10.1093/postmj/qgad006.
112. Cano Megías M, Alfaro Martínez JJ, Fernández de Bobadilla Pascual B, Bellerive G, García Manzanares Vázquez de Agredos Á, Álvarez de Frutos V. Malnutrition risk prevalence in hospitalised patients in Castilla-La Mancha: Ten years after the PREDyCES® study. *Endocrinol Diabetes Nutr (Engl Ed).* 2023;70:492–500. doi: 10.1016/j.endien.2023.07.003.
113. Bech CB, Svendsen JA, Knudsen AW, Munk T, Beck AM. The association between malnutrition and dehydration in older adults admitted to a geriatric unit: An observational study. *Clin Nutr ESPEN.* 2023;57:598–605. doi: 10.1016/j.clnesp.2023.08.011.
114. Bannert K, Sautter LF, Wiese ML et al. Analysis of ESPEN and GLIM algorithms reveals specific drivers for the diagnosis of malnutrition in patients with chronic gastrointestinal diseases. *Nutrition.* 2023;106:111887. doi: 10.1016/j.nut.2022.111887.
115. Alvarez-Larruy M, Tomsen N, Guanyabens N, Palomeras E, Clavé P, Nascimento W. Spontaneous Swallowing Frequency in Post-Stroke Patients with and Without Oropharyngeal Dysphagia: An Observational Study. *Dysphagia.* 2023;38:200–10. doi: 10.1007/s00455-022-10451-3.

116. Zhang Y, Zhang L, Gao X et al. Validation of the GLIM criteria for diagnosis of malnutrition and quality of life in patients with inflammatory bowel disease: A multicenter, prospective, observational study. *Clin Nutr.* 2022;41:1297–306. doi: 10.1016/j.clnu.2022.04.016.
117. Zhang Q, Yu S, Li Q, Zhang M, Meng L, Hu S. Preoperative Nutritional Status in Elderly Inpatients with Gastrointestinal Cancer and Its Linear Association with Frailty. *Nutr Cancer.* 2022;74:1376–87. doi: 10.1080/01635581.2021.1955284.
118. Yuksel B, Edeer AD. Evaluation of preoperative nutritional status of patients with gastrointestinal cancer using different nutritional screening tests. *Turk Geriatr Derg.* 2022;25(4):632–39. doi: 10.31086/tjgeri.2022.322.
119. Xu LB, Shi MM, Huang ZX et al. Impact of malnutrition diagnosed using Global Leadership Initiative on Malnutrition criteria on clinical outcomes of patients with gastric cancer. *JPEN J Parenter Enteral Nutr.* 2022;46:385–94. doi: 10.1002/jpen.2127.
120. Xie LL, Jiang JJ, Fu HB, Zhang WY, Yang L, Yang M. Malnutrition in Relation to Muscle Mass, Muscle Quality, and Muscle Strength in Hospitalized Older Adults. *J Am Med Dir Assoc.* 2022;23:722–28. doi: 10.1016/j.jamda.2021.11.025.
121. Xie B, Sun Y, Sun J, Deng T, Jin B, Gao J. Applicability of five nutritional screening tools in Chinese patients undergoing colorectal cancer surgery: a cross-sectional study. *BMJ Open.* 2022;12(5):e057765. doi: 10.1136/bmjopen-2021-057765.
122. Trollebø MA, Skeie E, Revheim I et al. Comparison of nutritional risk screening with NRS2002 and the GLIM diagnostic criteria for malnutrition in hospitalized patients. *Sci Rep.* 2022;12(1):19743. doi: 10.1038/s41598-022-23878-3.
123. Thomas J, Delaney C, Miller M. The ability of the Global Leadership Initiative on Malnutrition (GLIM) to diagnose protein-energy malnutrition in patients requiring vascular surgery: A validation study. *Br J Nutr.* 2023;129:49–53. doi: 10.1017/s0007114522000344.
124. Thiam CN, Mathavan S, Abdullah A, Chong EGM. Malnutrition among patients admitted to the subacute geriatric ward during the COVID-19 pandemic era: A cross-sectional study in a tertiary hospital in Malaysia. *Med J Malaysia.* 2022;77(3):313–19. doi: N/A.
125. Sun WJ, Li GC, Zhang J, Zhu J, Zhang Z. The role of nutritional assessment for predicting radiotherapy-induced adverse events in patients with gastric cancer. *Br J Radiol.* 2022;95(1130):20201004. doi: 10.1259/bjr.20201004.
126. Song MM, Zhang Q, Liu T et al. Efficacy of Global Leadership Initiative on Malnutrition as potential cachexia screening tool for patients with solid cancer. *Nutr J.* 2022;21(1):9. doi: 10.1186/s12937-022-00829-2.
127. Song D, Zhang L, Zhang Y et al. Risk factors for inpatient malnutrition and length of stay assessed by 'NutritionDay' in China. *Asia Pac J Clin Nutr.* 2022;31:561–69. doi: 10.6133/apjcn.202209\_31%283%29.0024.

128. Shabanpur M, Pourmahmoudi A, Nicolau J et al. The importance of nutritional status on clinical outcomes among both ICU and Non-ICU patients with COVID-19. *Clin Nutr ESPEN*. 2022;49:225–31. doi: 10.1016/j.clnesp.2022.04.016.
129. Santos MLDD, Leite LO, Lages ICF. Prevalence of malnutrition, according to the GLIM criteria, in patients who are the candidates for gastrointestinal tract surgery. *Arq Bras Cir Dig*. 2022;35:e1663. doi: 10.1590/0102-672020210002e1663.
130. Sánchez-Torralvo FJ, Contreras-Bolívar V, Ruiz-Vico M et al. Relationship between malnutrition and the presence of symptoms of anxiety and depression in hospitalized cancer patients. *Support Care Cancer*. 2022;30:1607–13. doi: 10.1007/s00520-021-06532-y.
131. Reiner J, Koch K, Woitalla J et al. Body impedance analysis to estimate malnutrition in inflammatory bowel disease patients - A cross-sectional study. *J Dig Dis*. 2022;23(12):687–94. doi: 10.1111/1751-2980.13155.
132. Quncuo C, Liang Y, Li QY et al. High Prevalence of Nutritional Risk Among Pulmonary Patients Living on the Tibetan Plateau. *Front Nutr*. 2022;9:872457. doi: 10.3389/fnut.2022.872457.
133. Polat O, Yuruyen M, Sonmezoz GB et al. Malnutrition risk frequency and independent risk factors associated with mortality in hospitalized elderly patients with COVID-19 in Turkey. *Asia Pac J Clin Nutr*. 2022;31:355–61. doi: 10.6133/apjcn.202209\_31(3).0003.
134. Oguri M, Ishii H, Yasuda K, Sumi T, Takahashi H, Murohara T. Combined prognostic value of malnutrition using GLIM criteria and renal insufficiency in elderly heart failure. *ESC Heart Fail*. 2022;9:704–11. doi: 10.1002/ehf2.13685.
135. Muresan BT, Núñez-Abad M, Artero A et al. Relation of Malnutrition and Nosocomial Infections in Cancer Patients in Hospital: An Observational Study. *J Nutr Metab*. 2022;2022:5232480. doi: 10.1155/2022/5232480.
136. Martins PM, Gomes TLN, Franco EP, Vieira LL, Pimentel GD. High neutrophil-to-lymphocyte ratio at intensive care unit admission is associated with nutrition risk in patients with COVID-19. *JPEN J Parenter Enteral Nutr*. 2022;46:1441–48. doi: 10.1002/jpen.2318.
137. Lang JL, Shao YA, Liao JH et al. Patient-Generated Subjective Global Assessment (PG-SGA) predicts length of hospital stay in lung adenocarcinoma patients. *Br J Nutr*. 2022;127:1543–48. doi: 10.1017/s0007114521003500.
138. Kobayashi H, Inoue T, Ogawa M, Abe T, Tanaka T, Kakiuchi M. Malnutrition diagnosed by the Global Leadership Initiative on Malnutrition criteria as a predictor of gait ability in patients with hip fracture. *Injury*. 2022;53:3394–400. doi: 10.1016/j.injury.2022.08.004.
139. Kaya AS, Bora S, Yetisyigit T. Evaluation of the Effect of Nutritional Status in Patients with Cancer Receiving Chemotherapy on Anthropometric Measurements and Quality of Life. *Nutr Cancer*. 2022;74:1994–2002. doi: 10.1080/01635581.2021.1984541.
140. Kasapoglu US, Gok A, Delen LA, Ozer AB. Comparison of nutritional risk status assessment tools in predicting 30-day survival in critically ill COVID-19 pneumonia patients. *Ann Saudi Med*. 2022;42:236–45. doi: 10.5144/0256-4947.2022.236.

141. Kaluzniak-Szymanowska A, Krzyminska-Siemaszko R, Deskur-Smielecka E, Lewandowicz M, Kaczmarek B, Wieczorowska-Tobis K. Malnutrition, Sarcopenia, and Malnutrition-Sarcopenia Syndrome in Older Adults with COPD. *Nutrients*. 2022;14(1):44. doi: 10.3390/nu14010044.
142. Jiang M, Sun MY, Zhang X, Li RJ. Nutritional status and its association with in-hospital major adverse cardiac events in patients with severe heart failure: a prospective study. *Nutr Hosp*. 2022;39:256–65. doi: 10.20960/nh.03846.
143. Huang SS, Niu Y, Liu XW, Gu ZY, Huang AY, Wu J. Characteristics of malnutrition according to Global Leadership Initiative on Malnutrition criteria in non-surgical patients with irritable bowel disease. *Nutrition*. 2022;94:111514. doi: 10.1016/j.nut.2021.111514.
144. Huang DD, Yu DY, Wang WB et al. Global leadership initiative in malnutrition (GLIM) criteria using hand-grip strength adequately predicts postoperative complications and long-term survival in patients underwent radical gastrectomy for gastric cancer. *Eur J Clin Nutr*. 2022;76:1323–31. doi: 10.1038/s41430-022-01109-2.
145. Hamdan MH, Badrasawi MM, Abu Alwafa RW. Nutrition and functional status among Palestinian cancer patients receiving chemotherapy. *J Taibah Univ Med Sci*. 2022;17(2):264–70. doi: 10.1016/j.jtumed.2021.11.006.
146. Gregoriano C, Voelkle M, Koch D et al. Association of Different Malnutrition Parameters and Clinical Outcomes among COVID-19 Patients: An Observational Study. *Nutrients*. 2022;14(16):3449. doi: 10.3390/nu14163449.
147. Gómez-Uranga A, Guzmán-Martínez J, Esteve-Atiénzar PJ et al. Nutritional and Functional Impact of Acute SARS-CoV-2 Infection in Hospitalized Patients. *J Clin Med*. 2022;11(9):2424. doi: 10.3390/jcm11092424.
148. Gao B, Chen W, Liu Y et al. Associations between nutrition risk scores and sarcopenia in gastrointestinal cancer patients: a cross-sectional study. *Support Care Cancer*. 2022;30:3269–77. doi: 10.1007/s00520-021-06729-1.
149. Efendioglu EM, Cigiloglu A, Turkbeyler IH. Malnutrition and Depressive Symptoms in Elderly Palliative Care Patients. *J Palliat Care*. 2022;37:503–09. doi: 10.1177/08258597211045500.
150. Dai LL, Li WL, Zheng DF, Wang WH, Xie HF, Ma JW. Prevalence and Management Recommendations for Disease-Related Malnutrition in Chronic Kidney Disease Patients with and without Diabetes. *Int J Endocrinol*. 2022;2022:4419486. doi: 10.1155/2022/4419486.
151. Dahl H, Warz SI, Welland NL, Arnesen I, Marti HP, Dierkes J. Factors associated with nutritional risk in patients receiving haemodialysis assessed by Nutritional Risk Screening 2002 (NRS2002). *J Ren Care*. 2022;48:112–18. doi: 10.1111/jorc.12374.
152. Chen LR, Liu CY, Cui HY et al. Malnutrition dynamics according to GLIM criteria in hospitalized elderly. *Asia Pac J Clin Nutr*. 2022;31:543–50. doi: 10.6133/apjcn.202209\_31(3).0022.
153. Chen J, Zhao C, Huang YZ et al. Malnutrition predicts poor outcomes in diabetic COVID-19 patients in Huangshi, Hubei. *J Biomed Res*. 2022;36:32–38. doi: 10.7555/jbr.35.20210083.

154. Carvalho J, Correia MA, Kanegusuku H et al. Association between the risk of malnutrition and functional capacity in patients with peripheral arterial disease: A cross-sectional study. *PLoS One*. 2022;17(9):e0273051. doi: 10.1371/journal.pone.0273051.
155. Can B, Senturk Durmus N, Olgun Yıldızeli S, Kocakaya D, Ilhan B, Tufan A. Nutrition risk assessed by Nutritional Risk Screening 2002 is associated with in-hospital mortality in older patients with COVID-19. *Nutr Clin Pract*. 2022;37:605–14. doi: 10.1002/ncp.10860.
156. Borges K, Artacho R, Jodar-Graus R, Molina-Montes E, Ruiz-Lopez MD. Calf Circumference, a Valuable Tool to Predict Sarcopenia in Older People Hospitalized with Hip Fracture. *Nutrients*. 2022;14(20):4255. doi: 10.3390/nu14204255.
157. Barros TA, Cruvel JMDS, Silva BDM et al. Agreement between nutritional risk screening tools and anthropometry in hospitalized pediatric patients. *Clin Nutr ESPEN*. 2022;47:227–32. doi: 10.1016/j.clnesp.2021.12.008.
158. Ahmadi S, Firoozi D, Dehghani M et al. Evaluation of Nutritional Status of Intensive Care Unit COVID-19 Patients Based on the Nutritional Risk Screening 2002 Score. *Int J Clin Pract*. 2022;2022:2448161. doi: 10.1155/2022/2448161.
159. Zhou LP, Yu DY, Ma BW et al. Feasibility of substituting handgrip strength for muscle mass as a constituent standard in the Global Leadership Initiative on Malnutrition for diagnosing malnutrition in patients with gastrointestinal cancers. *Nutrition*. 2021;84:111044. doi: 10.1016/j.nut.2020.111044.
160. Zhang Z, Wan Z, Zhu Y, Zhang L, Zhang L, Wan H. Prevalence of malnutrition comparing NRS2002, MUST, and PG-SGA with the GLIM criteria in adults with cancer: A multi-center study. *Nutrition*. 2021;83:111072. doi: 10.1016/j.nut.2020.111072.
161. Yu Y, Ye J, Chen M et al. Malnutrition Prolongs the Hospitalization of Patients with COVID-19 Infection: A Clinical Epidemiological Analysis. *J Nutr Health Aging*. 2021;25:369–73. doi: 10.1007/s12603-020-1541-y.
162. Tran TP, Nguyen LT, Hirose K et al. Malnutrition is associated with dysphagia in Vietnamese older adult inpatients. *Asia Pac J Clin Nutr*. 2021;30(4):588–94. doi: 10.6133/apjcn.202112\_30(4).0005.
163. Skeie E, Sygnestveit K, Nilsen RM, Harthug S, Koch AM, Tangvik RJ. Prevalence of patients "at risk of malnutrition" and nutritional routines among surgical and non-surgical patients at a large university hospital during the years 2008-2018. *Clin Nutr*. 2021;40:4738–44. doi: 10.1016/j.clnu.2021.05.029.
164. Santos IM, Mendes L, Carolino E, Santos CA. Nutritional Status, Functional Status, and Quality of Life - What is the Impact and Relationship on Cancer Patients? *Nutr Cancer*. 2021;73:2554–67. doi: 10.1080/01635581.2020.1839520.
165. Rosato E, Gigante A, Gasperini ML, Proietti L, Muscaritoli M. Assessing Malnutrition in Systemic Sclerosis With Global Leadership Initiative on Malnutrition and European Society of Clinical Nutrition and Metabolism Criteria. *JPEN J Parenter Enteral Nutr*. 2021;45:618–24. doi: 10.1002/jpen.1872.

166. Rivelrud M, Paur I, Sygnestveit K, Nilsen RM, Tangvik RJ. Nutritional treatment is associated with longer survival in patients with pancreatic disease and concomitant risk of malnutrition. *Clin Nutr.* 2021;40:2128–37. doi: 10.1016/j.clnu.2020.09.037.
167. Pironi L, Sasdelli AS, Ravaioli F et al. Malnutrition and nutritional therapy in patients with SARS-CoV-2 disease. *Clin Nutr.* 2021;40:1330–37. doi: 10.1016/j.clnu.2020.08.021.
168. Pimentel RFW, Moraes GCD, Barcelos SGC, Figueiredo PCMD, Das Merces MC. Evaluation of Nutritional Risk and Prevalence of Obesity in Patients with Covid-19 in A Reference Hospital in Salvador, Bahia, Brazil: A Cross-Sectional Study. *Int J Nutrol.* 2021;14(1):11–15. doi: 10.1055/s-0041-1728680.
169. Nozoe M, Yamamoto M, Masuya R, Inoue T, Kubo H, Shimada S. Prevalence of Malnutrition Diagnosed with GLIM Criteria and Association with Activities of Daily Living in Patients with Acute Stroke. *J Stroke Cerebrovasc Dis.* 2021;30(9):105989. doi: 10.1016/j.jstrokecerebrovasdis.2021.105989.
170. Nishioka S, Matsushita T, Yamanouchi A et al. Prevalence and Associated Factors of Coexistence of Malnutrition and Sarcopenia in Geriatric Rehabilitation. *Nutrients.* 2021;13(11):3745. doi: 10.3390/nu13113745.
171. Mitani A, Iwai T, Shichinohe T et al. The Combined Usage of the Global Leadership Initiative on Malnutrition Criteria and Controlling Nutrition Status Score in Acute Care Hospitals. *Ann Nutr Metab.* 2021;77:178–84. doi: 10.1159/000516994.
172. Marinho R, Pessoa A, Lopes M et al. Prevalence of Nutritional Risk at Admission in Internal Medicine Wards in Portugal: The Multicentre Cross-Sectional ANUMEDI Study. *Acta Med Port.* 2021;34:420–27. doi: 10.20344/amp.13182.
173. Liu Y, Wang LT, Li ST et al. Associations Between Blood Trace Element Levels and Nutritional Status in Maintenance Hemodialysis. *J Ren Nutr.* 2021;31:661–68. doi: 10.1053/j.jrn.2020.12.007.
174. Liu X, Cao J, Zhu Z et al. Nutritional risk and nutritional status in hospitalized older adults living with HIV in Shenzhen, China: a cross-sectional study. *BMC Infect Dis.* 2021;21(1):618. doi: 10.1186/s12879-021-06322-1.
175. Liu H, Song B, Jin J et al. Nutritional Risk, Health Outcomes, and Hospital Costs Among Chinese Immobile Older Inpatients: A National Study. *Front Nutr.* 2021;8:758657. doi: 10.3389/fnut.2021.758657.
176. Liu CY, Lu ZH, Li ZJ, Xu JY, Cui HY, Zhu MW. Influence of Malnutrition According to the GLIM Criteria on the Clinical Outcomes of Hospitalized Patients With Cancer. *Front Nutr.* 2021;8:774636. doi: 10.3389/fnut.2021.774636.
177. Liu AH, Cong JJ, Wang Q et al. Risk of Malnutrition Is Common in Patients with Coronavirus Disease 2019 (COVID-19) in Wuhan, China: A Cross-sectional Study. *J Nutr.* 2021;151:1591–96. doi: 10.1093/jn/nxab009.

178. Lee B, Han HS, Yoon YS. Impact of Preoperative Malnutrition on Postoperative Long-Term Outcomes of Patients With Pancreatic Head Cancer. *Ann Surg Open*. 2021;2(1):e047. doi: 10.1097/as9.0000000000000047.
179. Leandro-Merhi VA, de Aquino JLB, Bertelli HD, Ramos GG, Mendes ET, Mendonça JA. Factors associated with patient weight loss and prescribed diet during hospitalization. *Nutr Hosp*. 2021;38:749–57. doi: 10.20960/nh.03249.
180. Lai XX, Zhu HW, Du HD, Huo XP, Yu K. Nutritional status of Chinese oldest-old adults ( $\geq 80$  years of age): a cross-sectional study in Beijing. *Eur J Clin Nutr*. 2021;75:1040 – 46. doi: 10.1038/s41430-020-00826-w.
181. Hashemi ST, Kiaii BA, Mousavi S, Hosseinpour F. Evaluation of nutritional risk according to screening (NRS.2002) in multiple trauma patients admitted to ICU of Alzahra Hospital in Isfahan in 2018. *Pak J Med Health Sci*. 2021;15(6):1724–26. doi: 10.53350/pjmhs211561724.
182. Geng H, Wang SJ, Jin Y et al. Nutritional Status and Body Composition in Wilson Disease: A Cross-Sectional Study From China. *Front Nutr*. 2021;8:790520. doi: 10.3389/fnut.2021.790520.
183. Genest F, Rak D, Bätz E, Ott K, Seefried L. Sarcopenia and Malnutrition Screening in Female Osteoporosis Patients-A Cross-Sectional Study. *J Clin Med*. 2021;10(11):2344. doi: 10.3390/jcm10112344.
184. Celik ZM, Islamoglu AH, Sabuncular G, Toprak HS, Gunes FE. Evaluation of malnutrition risk of inpatients in a research and training hospital: A cross-sectional study. *Clin Nutr ESPEN*. 2021;41:261–67. doi: 10.1016/j.clnesp.2020.12.002.
185. Cao JJ, Xu HX, Li W et al. Nutritional assessment and risk factors associated to malnutrition in patients with esophageal cancer. *Curr Probl Cancer*. 2021;45(1):100638. doi: 10.1016/j.currproblcancer.2020.100638.
186. Balci C, Bolayir B, Eşme M et al. Comparison of the Efficacy of the Global Leadership Initiative on Malnutrition Criteria, Subjective Global Assessment, and Nutrition Risk Screening 2002 in Diagnosing Malnutrition and Predicting 5-Year Mortality in Patients Hospitalized for Acute Illnesses. *JPEN J Parenter Enteral Nutr*. 2021;45:1172–80. doi: 10.1002/jpen.2016.
187. Azman M, Teo DS, Yunus MRM, Yean KT. Mid-Upper Arm Circumference vs. Quadriceps Muscle Layer Thickness as a Surrogate Measure of Fat Free Mass Index for GLIM Criteria Assessment among Head and Neck Cancer Patients. *Int J Gerontol*. 2021;15(3):266–71. doi: 10.6890/ijge.202107\_15(3).0015.
188. Alikiaii B, Heidari Z, Fazeli A et al. Evaluation of the effectiveness of the Nutritional Risk Screening System 2002 (NRS-2002) in COVID-19 patients admitted to the intensive care unit. *Int J Clin Pract*. 2021;75(12):14934. doi: 10.1111/ijcp.14934.
189. Zhou X, Wu XN, Deng B, Huang LJ. Comparative Survey on Nutrition Risk and Nutrition Support Among Hospitalized General Surgery Patients Over a 7-Year Period. *JPEN J Parenter Enteral Nutr*. 2020;44:1468–74. doi: 10.1002/jpen.1784.

190. Zezelj SP, Malec D, Labinac DJ et al. Nutritional risk screening in gastroenterological patients at the rijeka university hospital centre. *Acta Clin Croat.* 2020;59(4):632–40. doi: 10.20471/acc.2020.59.04.09.
191. Xu JY, Zhu MW, Zhang H et al. A Cross-Sectional Study of GLIM-Defined Malnutrition Based on New Validated Calf Circumference Cut-off Values and Different Screening Tools in Hospitalised Patients over 70 Years Old. *J Nutr Health Aging.* 2020;24:832–38. doi: 10.1007/s12603-020-1386-4.
192. Skeie E, Tangvik RJ, Nymo LS, Harthug S, Lassen K, Viste A. Weight loss and BMI criteria in GLIM's definition of malnutrition is associated with postoperative complications following abdominal resections - Results from a National Quality Registry. *Clin Nutr.* 2020;39:1593–99. doi: 10.1016/j.clnu.2019.07.003.
193. Shimizu A, Maeda K, Honda T et al. Comparison between the Global Leadership Initiative on Malnutrition and the European Society for Clinical Nutrition and Metabolism definitions for the prevalence of malnutrition in geriatric rehabilitation care. *Geriatr Gerontol Int.* 2020;20(12):1221–27. doi: 10.1111/ggi.14072.
194. Sanz-París A, Martín-Palmero A, Gomez-Candela C et al. GLIM Criteria at Hospital Admission Predict 8-Year All-Cause Mortality in Elderly Patients With Type 2 Diabetes Mellitus: Results From VIDA Study. *JPEN J Parenter Enteral Nutr.* 2020;44:1492–500. doi: 10.1002/jpen.1781.
195. Matsumoto Y, Iwai K, Namikawa N et al. The relationship between existing nutritional indicators and Global Leadership Initiative on Malnutrition (GLIM) criteria: A one-institution cross-sectional analysis. *Clin Nutr.* 2020;39:3099–104. doi: 10.1016/j.clnu.2020.01.016.
196. Makwana NR, Ram RV, Khambhati S. Screening of malnutrition using the mini-nutritional assessment among hospitalized elderly patients in Western Gujarat. *Int J Health Allied Sci.* 2020;9(2):132–35. doi: 10.4103/ijhas.IJHAS\_102\_19.
197. Maeda K, Ishida Y, Nonogaki T, Mori N. Reference body mass index values and the prevalence of malnutrition according to the Global Leadership Initiative on Malnutrition criteria. *Clin Nutr.* 2020;39:180–84. doi: 10.1016/j.clnu.2019.01.011.
198. Ma J, Yang X, Cao J, Huang M, Jiang J, Wu R. A comparative study of three nutritional risk screening tools in surgical inpatients with laryngeal cancer. *Asia Pac J Clin Nutr.* 2020;29:227–33. doi: 10.6133/apjcn.202007\_29(2).0003.
199. Lueg G, Wirth R, Kwiatkowski J et al. Low Self-Perception of Malnutrition in Older Hospitalized Patients. *Clin Interv Aging.* 2020;15:2219–26. doi: 10.2147/cia.S278578.
200. Koroušić Seljak B, Mlakar Mastnak D, Mrevlje Ž, Veninšek G, Rotovnik Kozjek N. A multi-center survey on hospital malnutrition and cachexia in Slovenia. *Eur J Clin Nutr.* 2020;74:419–26. doi: 10.1038/s41430-019-0485-y.
201. Kaźmierczak-Siedlecka K, Skonieczna-Żydecka K, Folwarski M, Ruszkowski J, Świerblewski M, Makarewicz W. Influence of malnutrition stage according to GLIM 2019 criteria and SGA on the quality of life of patients with advanced cancer. *Nutr Hosp.* 2020;37:1179–85. doi: 10.20960/nh.03185.

202. Kaya AS, Pekcan G. Evaluation of preoperative nutritional status of patients with gastrointestinal cancer using different nutritional screening tests. *Prog Nutr.* 2020;22(3):e2020073. doi: 10.23751/pn.v22i3.9339.
203. Fukui S, Kawakami M, Otaka Y et al. Malnutrition among elderly patients with severe aortic stenosis. *Aging Clin Exp Res.* 2020;32:373–79. doi: 10.1007/s40520-019-01227-1.
204. Fernandes AC, Pessoa A, Vigário MA, Jager-Wittenaar H, Pinho J. Does malnutrition influence hospital reimbursement? A call for malnutrition diagnosis and coding. *Nutrition.* 2020;74:110750. doi: 10.1016/j.nut.2020.110750.
205. Dou L, Wang X, Cao Y, Hu A, Li L. Relationship between Postoperative Recovery and Nutrition Risk Screened by NRS 2002 and Nutrition Support Status in Patients with Gastrointestinal Cancer. *Nutr Cancer.* 2020;72:33–40. doi: 10.1080/01635581.2019.1612927.
206. De Groot LM, Lee G, Ackerie A, van der Meij BS. Malnutrition Screening and Assessment in the Cancer Care Ambulatory Setting: Mortality Predictability and Validity of the Patient-Generated Subjective Global Assessment Short form (PG-SGA SF) and the GLIM Criteria. *Nutrients.* 2020;12(8):2287. doi: 10.3390/nu12082287.
207. Chávez-Tostado M, Cervantes-Guevara G, López-Alvarado SE et al. Comparison of nutritional screening tools to assess nutritional risk and predict clinical outcomes in Mexican patients with digestive diseases. *BMC Gastroenterol.* 2020;20(1):79. doi: 10.1186/s12876-020-01214-1.
208. Burgos R, Joaquín C, Blay C, Vaqué C. Disease-related malnutrition in hospitalized chronic patients with complex needs. *Clin Nutr.* 2020;39:1447–53. doi: 10.1016/j.clnu.2019.06.006.
209. Boulhosa RSSB, Lourenco RP, Cortes DM, Oliveira LPM, Lyra AC, de Jesus RP. Comparison between criteria for diagnosing malnutrition in patients with advanced chronic liver disease: GLIM group proposal versus different nutritional screening tools. *J Hum Nutr Diet.* 2020;33:862–68. doi: 10.1111/jhn.12759.
210. Adly NN, Abd-El-Gawad WM, Abou-Hashem RM. Relationship between malnutrition and different fall risk assessment tools in a geriatric in-patient unit. *Aging Clin Exp Res.* 2020;32:1279–87. doi: 10.1007/s40520-019-01309-0.
211. Acehan S, Gulen M, Satar S et al. Evaluation of Nutrition Risk in Patients Over 65 Years of Age With Nontraumatic Acute Abdominal Syndrome. *Nutr Clin Pract.* 2020;35:1070–79. doi: 10.1002/ncp.10575.
212. Susetyowati, Budiningsari RD, Widodo P. Association of malnutrition screening tools with nutritional assessment parameters in hospitalized adult patients. *Pak J Nutr.* 2019;18(12):1145–52. doi: 10.3923/pjn.2019.1145.1152.
213. Ostrowska JM, Wegierek DS, Steinhagen AJ. Applying various criteria to assess the nutritional status among hospitalised patients aged 65 and over. *Prog Nutr.* 2019;21(1):113–19. doi: 10.23751/pn.v21i1.6952.
214. Ong SH, Chee WSS, Mageswary Lapchmanan L, Ong SN, Lua ZC, Yeo JXN. Validation of the Subjective Global Nutrition Assessment (SGNA) and Screening Tool for the Assessment of

- Malnutrition in Paediatrics (STAMP) to Identify Malnutrition in Hospitalized Malaysian Children. *J Trop Pediatr*. 2019;65(1):39–45. doi: 10.1093/tropej/fmy009.
215. Mishamandani ZJ, Norouzy A, Hashemian SM et al. Nutritional status of patients hospitalized in the intensive care unit: A comprehensive report from Iranian hospitals, 2018. *J Crit Care*. 2019;54:151–58. doi: 10.1016/j.jcrc.2019.08.001.
216. Miao JP, Quan XQ, Zhang CT et al. Comparison of two malnutrition risk screening tools with nutritional biochemical parameters, BMI and length of stay in Chinese geriatric inpatients: a multicenter, cross-sectional study. *BMJ Open*. 2019;9(2):e022993. doi: 10.1136/bmjopen-2018-022993.
217. Gutiérrez-de-Santiago JL, Aguilar-Valdez S, Casas-Robles ML, Garza-Veloz I, Ortega-Cisneros V, Martínez-Fierro ML. Screening of nutritional risk: assessment of predictive variables of nutritional risk in hospitalized patients in a second-level care center in Mexico. *Nutr Hosp*. 2019;36:626–32. doi: 10.20960/nh.2394.
218. Çoban E. Malnutrition Rate in Stroke Patients on Admission. *Sisli Etfal Hastan Tip Bul*. 2019;53(3):272–75. doi: 10.14744/semb.2018.81994.
219. Cao Q, Huang YH, Jiang M, Dai C. The prevalence and risk factors of psychological disorders, malnutrition and quality of life in IBD patients. *Scand J Gastroenterol*. 2019;54:1458–66. doi: 10.1080/00365521.2019.1697897.
220. Caliskan E, Dogan O. Synergistic effect of frailty and malnutrition on postoperative firstmonth mortality and delirium status among geriatric age group patients with hip fractures. *Turk Geriatr Derg*. 2019;22(2):140–49. doi: 10.31086/tjgeri.2019.87.
221. Aynaci G, Guksu Z. Nutritional screening and the impact of malnutrition on poor postoperative outcomes in gynecological oncology patients. *Prog Nutr*. 2019;21(1):127–34. doi: 10.23751/pn.v21i1.8044.
222. Rinninella E, Cintoni M, De Lorenzo A et al. Risk, prevalence, and impact of hospital malnutrition in a Tertiary Care Referral University Hospital: a cross-sectional study. *Intern Emerg Med*. 2018;13:689–97. doi: 10.1007/s11739-018-1884-0.
223. Kumar S, Umarji A, Bhangale K et al. Association of Nutritional Status with Outcome among Elderly Patients Admitted with Respiratory Diseases in a Critical Care Unit: A Cross-sectional Study. *J Clin Diagn Res*. 2025;19(8):OC44–OC47. doi: 10.7860/JCDR/2025/80353.21407.
224. Deng LM, Liu SP, Long JF et al. Prevalence of malnutrition and appropriateness of nutritional support in hospitalized patients: a GLIM-based study. *Front Nutr*. 2025;12:1667821. doi: 10.3389/fnut.2025.1667821.
225. WHO. Countries. [cited 2024/2/19]; Available from: <https://www.who.int/countries>
226. WHO. UN M49 – United Nations Statistics Division. 1999/1/1 [cited 2025/2/25]; Available from: <https://unstats.un.org/unsd/methodology/m49/>
227. Zhou J, Wang M, Wang H, Chi Q. Comparison of two nutrition assessment tools in surgical elderly inpatients in Northern China. *Nutr J*. 2015;14:68. doi: 10.1186/s12937-015-0054-8.

228. Ng DHL, Koh FHX, Yeong HEL et al. Nutrition Care after Hospital Discharge in Singapore: Evidence-Based Best-Practice Recommendations. *Nutrients*. 2023;15(21):4492. doi: 10.3390/nu15214492.
229. Huang XM, Lei Y, Xiao HX, Lian YY, Zhao Y. Consistency of "One-step" vs "Two-step" methods in malnutrition assessment according to global leadership initiative on malnutrition (GLIM) criteria among hospitalized patients with malignancy. *Parenter Enteral Nutr*. 2022;29(1):1–6. doi: 10.16151/j.1007-810x.2022.01.001.
230. Lian YY, Zhao Y. Effect of nutritional risk screening before assessment on GLIM criteria malnutrition diagnosis. *Electron J Metab Nutr Cancer*. 2021;8(5):525–29. doi: N/A.
231. Zeng YJ, Wang XL, Liu Y et al. An exploration of registered dietitian accreditation system development in China. *BMC Med Educ*. 2022;22(1):846. doi: 10.1186/s12909-022-03802-z.
232. Ishido K, Nakashima M, Poudel S et al. Implementation of the GLIM (Global Leadership Initiative on Malnutrition) Criteria in Gastrointestinal Oncology: A Narrative Review of Systemic Impact and the Role of Dietitians. *Cureus*. 2025;17(6):e86826. doi: 10.7759/cureus.86826.
233. Green SM, James EP. Barriers and facilitators to undertaking nutritional screening of patients: a systematic review. *J Hum Nutr Diet*. 2013;26:211–21. doi: 10.1111/jhn.12011.
234. Narayan SK, Gudivada KK, Krishna B. Assessment of Nutritional Status in the Critically Ill. *Indian J Crit Care Med*. 2020;24(Suppl 4):152–56. doi: 10.5005/jp-journals-10071-23617.
235. Guan X, Chen D, Xu Y. Clinical practice guidelines for nutritional assessment and monitoring of adult ICU patients in China. *J Intensiv Med*. 2024;4(2):137–59. doi: 10.1016/j.jointm.2023.12.002.
236. Ramaswamy T, DeWane MP, Dashti HS et al. Nine Myths about Enteral Feeding in Critically Ill Adults: An Expert Perspective. *Adv Nutr*. 2025;16(1):100345. doi: 10.1016/j.advnut.2024.100345.
237. McClave SA, Taylor BE, Martindale RG et al. Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *JPEN J Parenter Enteral Nutr*. 2016;40:159–211. doi: 10.1177/0148607115621863.
238. Chinese Society of Parenteral and Enteral Nutrition. Guideline for clinical application of parenteral and enteral nutrition in adults patients in China (2023 edition). *Zhonghua Yi Xue Za Zhi*. 2023;103(13):946–74. doi: 10.3760/cma.j.cn112137-20221116-02407.
239. ICDA. Dietitian-Nutritionists around the World Their education and their work (2021) 2023/11/1 [cited 2025/8/6]; Available from: <https://internationaldietetics.org/wp-content/uploads/2023/11/Education-and-Work-Report-2021.pdf>
240. Marcason W. What is ADA's staffing ratio for clinical dietitians? *J Am Diet Assoc*. 2006;106(11):1916. doi: 10.1016/j.jada.2006.09.025.
241. Varma S, Kapoor M, Patil M, Patel D, Timmanpyati S. Survey of Nutritional Screening Status as Well as Dietitian to Patient Ratio in Various Healthcare Setups in India. *J Nutr Res*. 2023;11(1):33–37. doi: 10.55289/jnutres/v11i1\_22.24.

242. Merlino Barr S, Hand RK, Fenton TR, Groh-Wargo S. Role of the neonatal registered dietitian nutritionist in Canada: A description of staffing and a comparison to practices in the United States. *Nutr Clin Pract*. 2024;39:1212–26. doi: 10.1002/ncp.11182.
243. Singer P, Robinson E, Raphaeli O. The future of artificial intelligence in clinical nutrition. *Curr Opin Clin Nutr Metab Care*. 2024;27:200–06. doi: 10.1097/MCO.0000000000000977.
244. Wang X, Yao K, Huang Z et al. An artificial intelligence malnutrition screening tool based on electronic medical records. *Clin Nutr ESPEN*. 2025;68:153–59. doi: 10.1016/j.clnesp.2025.03.178.
245. Long Z, Huang S, Zhang J et al. A Digital Smartphone-Based Self-administered Tool (R+ Dietitian) for Nutritional Risk Screening and Dietary Assessment in Hospitalized Patients With Cancer: Evaluation and Diagnostic Accuracy Study. *JMIR Form Res*. 2022;6(10):e40316. doi: 10.2196/40316.
246. Mancin S, Ferrara G, Lopane D et al. Artificial intelligence in the management of hospital malnutrition: A systematic review. *Clin Nutr ESPEN*. 2025;70:542–53. doi: 10.1016/j.clnesp.2025.10.002.
247. Janssen SM, Bouzembrak Y, Tekinerdogan B. Artificial Intelligence in Malnutrition: A Systematic Literature Review. *Adv Nutr*. 2024;15(9):100264. doi: 10.1016/j.advnut.2024.100264.
248. Wischmeyer PE, Bear DE, Berger MM et al. Personalized nutrition therapy in critical care: 10 expert recommendations. *Crit Care*. 2023;27(1):261. doi: 10.1186/s13054-023-04539-x.
249. Briassoulis G, Briassouli E. AI-Enabled Precision Nutrition in the ICU: A Narrative Review and Implementation Roadmap. *Nutrients*. 2025;18(1):110. doi: 10.3390/nu18010110.
250. Jiang ZM, Chen W, Zhu SN, Yao C. Survey on the Prevalence of Malnutrition (Inadequate Nutrition) and Nutritional Risk, and the Application of Nutritional Support in Tertiary Hospitals in Major Cities of Eastern, Central, and Western China. *Chin J Clin Nutr*. 2008;16:335–37. doi: 10.3881/j.issn.1008-5882.2008.06.002.
251. Sun H, Zhang L, Zhang P et al. A comprehensive nutritional survey of hospitalized patients: Results from nutritionDay 2016 in China. *PLoS One*. 2018;13(3):e0194312. doi: 10.1371/journal.pone.0194312.
252. Terrin N, Schmid CH, Lau J, Olkin I. Adjusting for publication bias in the presence of heterogeneity. *Stat Med*. 2003;22:2113–26. doi: 10.1002/sim.1461.
253. Seid AM, Fentahun N. Prevalence of malnutrition among old people in Africa: systematic review and meta-analysis. *BMJ Open*. 2022;12(11):e065197. doi: 10.1136/bmjopen-2022-065197.
254. Correia M, Perman MI, Waitzberg DL. Hospital malnutrition in Latin America: A systematic review. *Clin Nutr*. 2017;36:958–67. doi: 10.1016/j.clnu.2016.06.025.
255. Qi YM, Ma XD, Liu XD et al. Investigation and Analysis of Clinical Nutrition Medical Service and Quality Safety in Medical Institutions in China. *Chin Health Qual Manage*. 2021;28(11):34–40. doi: 10.13912/j.cnki.chqm.2021.28.11.08.
256. Portugal Office of the Secretary of State for the Promotion of Health. Order No. 9984/2023 on the implementation and strengthening of systematic nutritional risk screening in all levels of care of the

National Health Service. 2023/9/27 [cited 2026/3/9]; Available from: <https://european-nutrition.org/good-practices/portugal-is-one-step-closer-to-mandatory-nutritional-e-screening/>

257. Mareschal J, Altwegg J, Berthet D et al. Prescription and indication for oral nutritional supplements in a Swiss university hospital: a prospective survey. *Swiss Med Wkly*. 2017;147:w14475. doi: 10.4414/smw.2017.14475.

258. National Healthcare Security Administration, Ministry of Human Resources and Social Security. Notice on Issuing the National Essential Medicines List for Basic Medical Insurance, Maternity Insurance, and Work Injury Insurance, and the Innovative Medicines List for Commercial Health Insurance (2025). 2025/12/7 [cited 2026/5/17]; Available from: [https://www.nhsa.gov.cn/art/2025/12/7/art\\_104\\_18970.html](https://www.nhsa.gov.cn/art/2025/12/7/art_104_18970.html).

Not Proof Read

**Table 1.** Pooled prevalence estimates of nutritional risk or malnutrition by screening tool and assessment timepoint

Tool	Target Population	Type / meaning	Time	Number of studies included / I <sup>2</sup>	Pooled prevalence (95% CI)	Publication bias (Peters' <i>p</i> ) <sup>†</sup>	Adjusted prevalence (Trim-and-fill / Bayesian)
NRS2002 ≥ 3	Adults (19–90y)	Screening / at nutritional risk	Discharge	3 / 94.6%	0.19 (0.09, 0.40)	– <sup>‡</sup> < 0.001	– 0.22 (0.18, 0.27) / 0.40 (95% CrI, 0.36, 0.45)
			Admission	90 / 99.9%	0.41 (0.35, 0.49)		
MNA-SF ≤ 11	Elderly (≥60y)	Screening / at risk of malnutrition	Admission	23 / 99.9%	0.53 (0.40, 0.71)	0.141	–
			Admission	30 / 99.3%	0.62 (0.57, 0.67)	0.977	–
STAMP ≥ 2	Children (2–17y)	Screening / at medium and high risk of malnutrition	Admission	6 / 98.5%	0.55 (0.44, 0.68)	–	–
GLIM	Adults	Diagnosis / malnutrition	Admission	80 / 99.6%	0.39 (0.35, 0.44)	< 0.001	0.26 (0.23, 0.30) / 0.39 (95% CrI, 0.35, 0.43)
One-Step GLIM	Adults	Diagnosis / malnutrition	Admission	44 / 99.1%	0.42 (0.37, 0.48)	0.021	0.35 (0.30, 0.40) / 0.42 (95% CrI, 0.37, 0.47)
Two-Step GLIM	Elderly (≥60y)	Diagnosis after NRS2002 malnutrition	Admission	5 / 99.5%	0.38 (0.21, 0.72)	–	–
	Adults (19–90y)			18 / 99.7%	0.34 (0.26, 0.46)	0.016	0.20 (0.15, 0.26) / 0.34 (95%CrI, 0.27, 0.43)
	Elderly (≥60y)			6 / 97.8%	0.28 (0.21, 0.37)	–	–
	Elderly (≥60y)	Diagnosis after MNA-SF / malnutrition	Admission	7 / 99.3%	0.49 (0.35, 0.68)	–	–

CI: confidence interval; CrI: credible interval; I<sup>2</sup>: measure of heterogeneity

<sup>†</sup>p values in the Publication Bias column are based on Peters' test

<sup>‡</sup>–, not assessed/not applicable.




Figure 1. PRISMA 2020 flow diagram of study selection




Figure 2. Summary forest plot of pooled prevalence of NRS2002 at admission and discharge (details in Supplementary Figure 1)




Figure 3. Forest plot of pooled prevalence of MNA-SF at admission




Figure 4. Forest plot of pooled prevalence of STAMP at admission




Figure 5. Summary forest plot of pooled prevalence of One-step GLIM and Two-step GLIM at admission (details in Supplementary Figure 2)




Figure 6. Summary of pooled prevalence at admission across countries