

Original Article

The impact of daily use of an enteral feeding checklist on clinical outcomes in shock patients: a retrospective cohort study

Zhi Mao MD¹, Qing Yu MD², Chao Liu MD³, Pan Hu MM¹, Xin Hu MM¹, Liang Pan MD¹, Hongjun Kang MD¹, Feihu Zhou MD, PhD^{1,3,4}

¹Department of Critical Care Medicine, Chinese People's Liberation Army General Hospital, Beijing, People's Republic of China

²Chifeng Municipal Hospital, Chifeng city of the Inner Mongolia Autonomous Region, People's Republic of China

³National Clinical Research Center for Kidney Diseases, Chinese People's Liberation Army General Hospital, Beijing, People's Republic of China

⁴National Clinical Research Center for Geriatric Diseases, Chinese People's Liberation Army General Hospital, Beijing, People's Republic of China

Background and Objectives: The optimal delivery of enteral nutrition in shock patients has an important prognostic clinical value; thus, checklists for standardizing enteral nutrition should be developed. This study examined whether the use of an enteral feeding checklist can improve enteral nutrition in shock patients. **Methods and Study Design:** A retrospective cohort study was conducted. A multidisciplinary working group developed an enteral feeding checklist. Information on patients' demographics, checklist items, and clinical outcomes was collected. **Results:** In total, 148 patients were included. The checklist was used for 35 patients but not for the remaining 113 patients. Patients in the checklist group received enteral nutrition earlier (2.6 vs 4.6 days, $p=0.017$) and had a lower mechanical ventilation rate (62.9% vs 85.0%, $p=0.004$). The checklist group had shorter intensive care unit stay (mean 17.3 vs 25.7 days, $p=0.043$). No significant differences were observed in 28- and 90-day mortality, mechanical ventilation duration, and intolerance to enteral nutrition. **Conclusions:** The use of an enteral feeding checklist in shock patients was associated with earlier enteral nutrition delivery and decreased intensive care unit stay.

Key Words: enteral feeding, checklist, shock patients, retrospective cohort

INTRODUCTION

Shock patients often have inadequate enteral intake and hypocalcemia, which increase the risks of malnutrition and mortality.^{1,2} Malnutrition and underfeeding may exist in more than 40% of critically ill patients.³ Providing appropriate nutritional support to critically ill patients can prevent malnutrition and improve clinical outcomes. Nutrition delivery methods for critical patients mainly include parenteral nutrition (PN) and enteral nutrition (EN). Being widely accepted, EN is preferred over PN. However, optimized EN delivery is delayed due to many reasons, such as physicians' delay in making decisions, underestimated energy demand, interrupted feeding, recent abdominal surgery, hemodynamic instability, and gastrointestinal abnormality.⁴⁻⁶ Several clinical practice guidelines have standardized EN delivery by involving a battery of interventions and procedures for nutrition therapy.⁷⁻⁹ In addition, enteral feeding protocols were introduced to improve clinical outcomes in many studies. However, although EN feeding protocols could increase the proportion of EN feeding, they failed to reduce mor-

tality, the incidence of nosocomial infection, and the duration of mechanical ventilation (MV).^{2,4}

Checklists help in preventing omission errors and are useful while performing tasks, ranging from simple shopping to flying an airplane.^{10,11} The use of checklists has also extended to intensive care units; for example, checklists are used while transferring critically ill patients,¹² reducing the incidence of catheter-related blood stream infection,¹³ maintaining hand hygiene compliance,¹⁴ reducing extubation failure,¹⁵ scheduling physical rehabilitation consultations, and conducting daily rounds. However, studies exploring the use of checklists for improving

Corresponding Author: Prof Feihu Zhou, Chinese People's Liberation Army General Hospital, 28 Fu-Xing Road, Beijing 100853, People's Republic of China.

Tel: 86-10-66938148; Fax: 86-10-88219862

Email: feihuzhou301@126.com

Manuscript received 31 January 2019. Initial review completed 20 February 2019. Revision accepted 10 March 2019.

doi:

EN in shock patients are limited. Therefore, the present study investigated the effect of using an EN checklist on shock patients.

METHODS

Study design

In this single-center retrospective cohort study, a before–after study design was used. The study protocol was approved by the Ethics Committee of PLA General Hospital (approval No. 2017-054-01). This study was conducted in accordance with the STROBE checklist.

Setting

The Department of Critical Care Medicine of PLA General Hospital is a multidisciplinary 20-bed unit and had an average of 800 annual admissions in December 1, 2015–Jun 30, 2017 and Jul 1, 2017–February 30, 2018.

Participants

All shock patients (age ≥ 18 years) who were admitted to the intensive care unit (ICU) and received EN and vasopressors were potentially eligible. We excluded patients who had contraindications to EN, including bowel obstruction, massive gastrointestinal bleeding, acute phase of severe pancreatitis, and post gastrointestinal operation; had received EN in the previous week; underwent percutaneous endoscopic jejunostomy; and had an estimated lifespan of < 24 h.

Use of a checklist

We designed a preliminary form that included patient's demographic data, studied practices, and other items for EN therapy assessment. Five staff physicians, who were responsible for the ICU and the implementation of the checklist, reviewed the draft form. We conducted a pilot test for this preliminary form between June 1, 2017, and July 30, 2017. Modifications were made in the design of the preliminary form according to the findings of the pilot test. The final checklist was confirmed by five staff physicians before its implementation (Figure 1).

Items in the checklist included hemodynamic data, acute gastrointestinal injury (AGI) score, nutritional risk assessment, method of nutrition, assessment of the aspiration risk, feeding route of EN, the EN product, caloric density of the EN product (kcal/mL), speed of EN delivery, assessment of EN tolerability, adjustment of EN project, and total dose of EN delivery (Figure 1). The calculation of the aspiration risk score was modified from a previous study (Supplement table 1).¹⁶

Variables and outcomes

Variables included patient demographics, Acute Physiology and Chronic Health Evaluation (APACHE) II score, and Sequential Organ Failure Assessment (SOFA) score over the first 24 h of admission to the ICU. The primary outcome was 28-day mortality. Secondary outcomes were 90-day mortality, length of stay in the ICU, duration of MV, and intolerance to EN feeding.

Statistical methods

All patients were divided into checklist and control groups. We reported quantitative variables as means with

standard deviations (SDs) for outcome measures with a normal distribution and as interquartile ranges for those with an abnormal distribution and categorical variables as rates. We performed the univariate analysis to determine statistical differences in these variables between the two groups. The univariate analysis included Student's *t* test, chi-square test, and Fisher's exact test, and $p < 0.05$ was considered significant. Multivariable logistic regression models were used to estimate the role of checklist use in clinical outcomes, and adjusted ORs with 95% CIs were calculated to estimate risks. Statistical analyses were performed using Empower (R) (<http://www.empowerstats.com>; X&Y Solutions Inc, Boston, Mass) and R software, version 3.1.2 (<http://www.R-project.org>).

RESULTS

A total of 148 patients were included in this study. The checklist was used in 35 (23.65 %) patients. The checklist assessment was not performed in the remaining 113 (76.35%) patients (Figure 2). The demographics of patients are listed in Table 1. No significant differences in sex, age, BMI, APACHE II score, SOFA score, and AGI score were found between the two groups in the univariate analysis (Table 1).

Compared with the control group, the checklist group received EN earlier, with a mean of 2.6 days versus 4.6 days ($p=0.017$), and had a lower rate of MV (62.9% vs 85.0%, $p=0.004$; Table 1).

The results of the univariate analysis of clinical outcomes are shown in Table 2. No significant difference was found between the two groups in 28-day mortality (20.0% vs 23.9%, $p=0.632$), 90-day mortality (25.7% vs 31.9%, $p=0.490$), duration of MV (mean 13.4 days vs 16.6 days; $p=0.395$), and intolerance to EN (17.1% vs 23.0%, $p=0.461$). However, the checklist group had shorter ICU stay (mean 17.3 vs. 25.7 days; $p=0.043$). The result of the multivariable logistic regression model also showed no significant difference in 28-day mortality, 90-day mortality, duration of MV, and intolerance to EN between the two groups. However, the checklist group still had significantly shorter ICU stay after adjustment for confounders (Table 3).

DISCUSSION

The results of this retrospective cohort study demonstrated that the EN checklist could reduce the length of ICU stay. However, a decrease was not observed in 28- and 90-day mortality or the durations of MV and hospitalization. In terms of baseline differences, only the MV rate differed. Enteral feeding was provided earlier in the checklist group. Other studies^{17,18} on EN protocols have also reported similar findings that support early EN.

The idea of using a checklist to prevent mistakes in an ICU was inspired by the inventory checks used in aviation. The daily management of ICU patients has many similarities to that of the aviation industry. For example, both industries involve timely assessment and management of complex multisystem objects, and minor errors may lead to serious adverse consequences. We referred to previous studies on the use of checklists in an ICU and adopted pre- and post-control study methods. The check-

Checklist of Nutrition Management In Criti

Checklist of Nutrition Management In Criti															
Data Name _____		ID _____		Date Of ICU Admission (Y/M/D) ____/____/____					The Dignosis Of ICU Admission _____						
Wight _____ kg		Energy Object _____ kcal/d			Protein Object _____ g/d										
Hemodynamic Stabilization (MAP>65mmHg, and The Dose of Vasopressor Is Decreased)	D1 <input type="checkbox"/> Yes <input type="checkbox"/> No					D2 <input type="checkbox"/> Yes <input type="checkbox"/> No					D3 <input type="checkbox"/> Yes <input type="checkbox"/> No				
	Dopamine	Norepinephrine	Epinephrine	Phenylephrine	Other Vasopressor	Dopamine	Norepinephrine	Epinephrine	Phenylephrine	Other Vasopressor	Dopamine	Norepinephrine	Epinephrine	Phenylephrine	Other Vasopressor
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Assessment of Gastrointestinal Function	<input type="checkbox"/> Normal/Mild Injury (AGI ≤ I grade)					<input type="checkbox"/> Normal/Mild Injury (AGI ≤ I grade)					<input type="checkbox"/> Normal/Mild Injury (AGI ≤ I grade)				
	<input type="checkbox"/> Moderate To Severe Injury (AGI II-III grade)					<input type="checkbox"/> Moderate To Severe Injury (AGI II-III grade)					<input type="checkbox"/> Moderate To Severe Injury (AGI II-III grade)				
	<input type="checkbox"/> Failure (AGI IV grade)					<input type="checkbox"/> Failure (AGI IV grade)					<input type="checkbox"/> Failure (AGI IV grade)				
EN intolerance risk	<input type="checkbox"/> High <input type="checkbox"/> Low <input type="checkbox"/> Not Evaluated					<input type="checkbox"/> High <input type="checkbox"/> Low <input type="checkbox"/> Not Evaluated					<input type="checkbox"/> High <input type="checkbox"/> Low <input type="checkbox"/> Not Evaluated				
Method of Nutrition	<input type="checkbox"/> EN <input type="checkbox"/> PN					<input type="checkbox"/> EN <input type="checkbox"/> PN					<input type="checkbox"/> EN <input type="checkbox"/> PN				
Assessment of Aspiration Risk	<input type="checkbox"/> High <input type="checkbox"/> Low <input type="checkbox"/> Not Evaluated					<input type="checkbox"/> High <input type="checkbox"/> Low <input type="checkbox"/> Not Evaluated					<input type="checkbox"/> High <input type="checkbox"/> Low <input type="checkbox"/> Not Evaluated				
Feeding Route of EN	<input type="checkbox"/> Enteral <input type="checkbox"/> Parenteral					<input type="checkbox"/> Enteral <input type="checkbox"/> Parenteral					<input type="checkbox"/> Enteral <input type="checkbox"/> Parenteral				
Name of The EN product															
Caloric Density of The EN product (kcal/ml)															
The Speed of Enteral Nutrition Delivered	_____ ml/h					_____ ml/h					_____ ml/h				
Assessment of EN Tolerability															
First Assessment of EN Tolerability	____ Score <input type="checkbox"/> Not Evaluated					____ Score <input type="checkbox"/> Not Evaluated					____ Score <input type="checkbox"/> Not Evaluated				
Adjustment of EN Project	<input type="checkbox"/> Increase <input type="checkbox"/> Invariability					<input type="checkbox"/> Increase <input type="checkbox"/> Invariability					<input type="checkbox"/> Increase <input type="checkbox"/> Invariability				
	<input type="checkbox"/> Decrease <input type="checkbox"/> Pause					<input type="checkbox"/> Decrease <input type="checkbox"/> Pause					<input type="checkbox"/> Decrease <input type="checkbox"/> Pause				
	____ Score <input type="checkbox"/> Not Evaluated					____ Score <input type="checkbox"/> Not Evaluated					____ Score <input type="checkbox"/> Not Evaluated				
	<input type="checkbox"/> Increase <input type="checkbox"/> Invariability					<input type="checkbox"/> Increase <input type="checkbox"/> Invariability					<input type="checkbox"/> Increase <input type="checkbox"/> Invariability				
	<input type="checkbox"/> Decrease <input type="checkbox"/> Pause					<input type="checkbox"/> Decrease <input type="checkbox"/> Pause					<input type="checkbox"/> Decrease <input type="checkbox"/> Pause				
The Total Dose of Enteral Nutrition Delivered (ml)															

Figure 1. Checklist of nutrition management in critical care medicine. Energy object: 25-30 kcal / kg (standard body weight)/d; (standard body weight (female) = Height (cm) - 105; (standard body weight (male) = height (cm)-110). Protein object: 1.2-2.0 g/kg (standard body weight) / d. Time record: time of admission to ICU is D0, the next day from beginning 8:00 am is D1, et al. Time of filling the form: 10:00 and 23:00. *EN intolerance high risk is defined as a patient having gastric residue >250 ml, vomiting, abdominal plain film or abdominal CT positive, intestinal ischemia or perforation.

ritical Care Medicine														
a _____										Height _____ cm				
D4					D5					D6				
<input type="checkbox"/> Yes <input type="checkbox"/> No					<input type="checkbox"/> Yes <input type="checkbox"/> No					<input type="checkbox"/> Yes <input type="checkbox"/> No				
Dopamine	Norepinephrine	Epinephrine	Phenylephrine	Other Vasopressor	Dopamine	Norepinephrine	Epinephrine	Phenylephrine	Other Vasopressor	Dopamine	Norepinephrine	Epinephrine	Phenylephrine	Other Vasopressor
<input type="checkbox"/> Normal/Mild Injury (AGI ≤ I grade)					<input type="checkbox"/> Normal/Mild Injury (AGI ≤ I grade)					<input type="checkbox"/> Normal/Mild Injury (AGI ≤ I grade)				
<input type="checkbox"/> Moderate to Severe Injury (AGI II-III grade)					<input type="checkbox"/> Moderate To Severe Injury (AGI II-III grade)					<input type="checkbox"/> Moderate To Severe Injury (AGI II-III grade)				
<input type="checkbox"/> Failure (AGI IV grade)					<input type="checkbox"/> Failure (AGI IV grade)					<input type="checkbox"/> Failure (AGI IV grade)				
<input type="checkbox"/> High <input type="checkbox"/> Low <input type="checkbox"/> Not Evaluated					<input type="checkbox"/> High <input type="checkbox"/> Low <input type="checkbox"/> Not Evaluated					<input type="checkbox"/> High <input type="checkbox"/> Low <input type="checkbox"/> Not Evaluated				
<input type="checkbox"/> EN <input type="checkbox"/> PN					<input type="checkbox"/> EN <input type="checkbox"/> PN					<input type="checkbox"/> EN <input type="checkbox"/> PN				
<input type="checkbox"/> High <input type="checkbox"/> Low <input type="checkbox"/> Not Evaluated					<input type="checkbox"/> High <input type="checkbox"/> Low <input type="checkbox"/> Not Evaluated					<input type="checkbox"/> High <input type="checkbox"/> Low <input type="checkbox"/> Not Evaluated				
<input type="checkbox"/> Enteral <input type="checkbox"/> Parenteral					<input type="checkbox"/> Enteral <input type="checkbox"/> Parenteral					<input type="checkbox"/> Enteral <input type="checkbox"/> Parenteral				
_____ ml/h					_____ ml/h					_____ ml/h				
_____ Score <input type="checkbox"/> Not Evaluated					_____ Score <input type="checkbox"/> Not Evaluated					_____ Score <input type="checkbox"/> Not Evaluated				
<input type="checkbox"/> Increase <input type="checkbox"/> Invariability					<input type="checkbox"/> Increase <input type="checkbox"/> Invariability					<input type="checkbox"/> Increase <input type="checkbox"/> Invariability				
<input type="checkbox"/> Decrease <input type="checkbox"/> Pause					<input type="checkbox"/> Decrease <input type="checkbox"/> Pause					<input type="checkbox"/> Decrease <input type="checkbox"/> Pause				
_____ Score <input type="checkbox"/> Not Evaluated					_____ Score <input type="checkbox"/> Not Evaluated					_____ Score <input type="checkbox"/> Not Evaluated				
<input type="checkbox"/> Increase <input type="checkbox"/> Invariability					<input type="checkbox"/> Increase <input type="checkbox"/> Invariability					<input type="checkbox"/> Increase <input type="checkbox"/> Invariability				
<input type="checkbox"/> Decrease <input type="checkbox"/> Pause					<input type="checkbox"/> Decrease <input type="checkbox"/> Pause					<input type="checkbox"/> Decrease <input type="checkbox"/> Pause				

Figure 1. Checklist of nutrition management in critical care medicine (cont.). Energy object: 25-30 kcal / kg (standard body weight)/d; (standard body weight (female) = Height (cm) - 105; (standard body weight (male) = height (cm)-110). Protein object: 1.2-2.0 g/kg (standard body weight) / d. Time record: time of admission to ICU is D0, the next day from beginning 8:00 am is D1, et al. Time of filling the form: 10:00 and 23:00. *EN intolerance high risk is defined as a patient having gastric residue >250 ml, vomiting, abdominal plain film or abdominal CT positive, intestinal ischemia or perforation.

Table 1. Baseline characteristics in each group

	Checklist Group	Control Group	p-value
N	35	113	
Sex			0.736
Female	11 (31.4%)	39 (34.5%)	
Male	24 (68.6%)	74 (65.5%)	
Age	59.4±18.8	60.5±19.3	0.764
Height (cm)	166.2±7.3	168.5±7.6	0.117
Weight (kg)	64.3±11.4	64.5±11.2	0.921
BMI	23.7±3.6	23.0±3.7	0.273
CRRT			0.255
No	24 (72.7%)	70 (61.9%)	
Yes	9 (27.3%)	43 (38.1%)	
MV			0.004
No	13 (37.1%)	17 (15.0%)	
Yes	22 (62.9%)	96 (85.0%)	
MAP (mmHg)	76.3±7.5	77.1±8.7	0.613
Abdominal diseases			0.407
No	27 (77.1%)	79 (69.9%)	
Yes	8 (22.9%)	34 (30.1%)	
EN start time (d)	2.6±2.3	4.6±4.7	0.017
Feeding way			0.050
No	10 (28.6%)	16 (14.2%)	
Yes	25 (71.4%)	97 (85.8%)	
EN total (ml)	500 (250-500) †	500 (250-500) †	0.541
EN Speed (ml/h)	25 (25-50) †	25 (20-50) †	0.309
Creatinine (mmol/L)	89 (73.6-98) †	86 (64-156) †	0.266
IL-6 (pg/ml)	108 (68-183) †	80.8 (30.0-227) †	0.144
Lactate (mmol/L)	1.8±1.1	2.0±1.7	0.498
SOFA	8.5±3.9	9.1±3.8	0.404
APACHE	18.7±8.0	17.7±7.7	0.504
AGI			0.335
1	25 (71.4%)	80 (70.8%)	
2	5 (14.3%)	21 (18.6%)	
3	3 (8.6%)	11 (9.7%)	
4	2 (5.7%)	1 (0.9%)	

AGI: acute gastrointestinal abbreviated injury; APACHE: acute physical and chronic health assessment; CRRT: continuous renal replacement therapy; EN: enteral nutrition; MV: mechanical ventilation; SOFA: sequential organ failure assessment.

† Non-parametric test, median(interquartile range).


**Figure 2.** Patients included in study cohorts.

Table 2. The comparison of characteristics and outcomes between checklist group and control group

	Checklist Group	Control Group	p-value
N	35	113	
Hospital stay (d)	33.7±22.5	39.0±25.2	0.275
ICU stay (d)	17.3±18.1	25.7±21.9	0.043
Duration of MV(d)	13.4±16.6	16.6±20.0	0.395
28-days mortality	7 (20.0%)	27 (23.9%)	0.632
90-days mortality	9 (25.7%)	36 (31.9%)	0.490
Intolerance of EN feeding	6 (17.1%)	26 (23.0%)	0.461

ICU: intensive care unit; MV: mechanical ventilation.

Table 3. The comparison of characteristics and outcomes between Checklist group and control group in multivariate analysis

Exposure	Non-adjusted	Adjust I	Adjust II
ICU stay (d)			
Control group	0	0	0
Checklist group	-8.3 (-16.3, -0.3) 0.043	-5.2 (-10.0, -0.5) 0.033	-5.0 (-9.8, -0.1) 0.048
Hospital stay (d)			
Control group	0	0	0
Checklist group	-5.2 (-14.5, 4.1) 0.275	-3.0 (-11.1, 5.1) 0.468	-2.7 (-10.9, 5.6) 0.527
Intolerance of EN feeding			
Control group	1.0	1.0	1.0
Checklist group	0.7 (0.3, 1.8) 0.463	1.0 (0.4, 2.9) 0.954	1.0 (0.3, 2.8) 0.952
28-days mortality			
Control group	1.0	1.0	1.0
Checklist group	0.8 (0.3, 2.0) 0.633	1.5(0.5, 4.1) 0.483	1.0 (0.3, 3.2) 0.959
90-days mortality			
Control group	1.0	1.0	1.0
Checklist group	0.7 (0.3, 1.7) 0.491	1.4 (0.5, 3.5) 0.520	0.9 (0.3, 2.7) 0.793

APACHE: acute physical and chronic health assessment; EN: enteral nutrition; ICU: intensive care unit; MV: mechanical ventilation. Data in the table: OR (95% CI) p-value.

Model I: Non-adjusted.

Model II adjust for: duration of MV, EN start time, MV, Abdominal diseases.

Model III adjust for: duration of MV, EN start time, MV, APACHE II Abdominal diseases.

list was based on studies focusing on hemodynamic stabilization, gastrointestinal function assessment, AGI, aspiration risk assessment, nutrition initiation, speed, risk assessment, and intolerance.

The hemodynamic status of shock patients was assessed in our checklist. Unstable hemodynamic status and vasopressors often cause gastrointestinal dysfunction and intolerance to feeding.¹⁹ Mancl et al²⁰ reported that the tolerance rate of EN in shock patients who received 12.5 µg•min⁻¹ of norepinephrine was 75%. Therefore, the dose of vasopressors was also recorded in our checklist.

The AGI grading system was proposed by the European Society of Intensive Care Medicine (ESICM) in 2012. The AGI grading system has been associated with the severity of gastrointestinal dysfunction and mortality.^{21,22} The checklist included the AGI grading system as an item.

Intolerance to feeding, which is a major cause of insufficient nutrition, is associated with mortality.^{20,23} The assessment of feeding intolerance included nausea, vomiting, constipation, diarrhea, gastrointestinal hemorrhage, and positive abdominal imaging findings. In addition, our EN checklist encourages staff to promptly treat feeding intolerance by adjusting the EN speed after the first assessment of EN tolerability.

Many studies have been conducted on the use of checklists in an ICU; however, no study has evaluated the benefits of using an EN checklist in ICU patients. Most of the studies on the implementation and management of EN have used an EN protocol. Li et al designed an EN protocol for critically ill patients and conducted a before and after study. They reported that EN could not reduce the mortality rate and MV duration.² Kim et al. also reported that the EN protocol could begin enteral nutrition early but did not affect mortality.⁴ Wikjord et al demonstrated that the EN protocol increased the proportion of early EN but did not affect clinical outcomes, such as length of ICU stay.³ Volume-based EN adjusts feeding depending on the hourly situation. However, no significant improvement was noted in indicators such as mortality, length of ICU stay, and MV duration.²⁴

Previous studies on the use of checklists in an ICU have involved nutrition management. Weiss et al²⁵ designed a daily checklist that contained an item on nutrition; that is, the percentage of nutrition goals achieved. The results of this study can be applied to a single-centre MICU to improve medical quality and reduce disease severity, mortality, and hospitalization duration. Centofanti et al²⁶ conducted a mixed methods study to investigate the effect of a daily goals checklist on rounds. In

their checklist, EN assessment was included as one of the items. They found that the daily goals checklist enhanced patient safety. Brunsveld-Reinders et al¹² also introduced EN evaluation in the post-transfer part of their transport checklist. However, few studies have performed the nutrition assessment of shock patients.

This study has some limitations. First, because this is a retrospective cohort study, all limitations and bias of the retrospective cohort study were unavoidable. The checklist group had a higher MV rate and earlier EN start; this might have confounded the causality. However, the multivariate analysis confirmed the results. Second, the generalizability of this single-center study may be limited. Third, because this is a pilot study, the sample size was small. Thus, larger studies with prospective randomized controlled trial design are needed.

Conclusion

The implementation of the EN checklist for shock patients in an ICU could reduce the length of ICU stay. However, the use of the EN checklist did not improve mortality, MV duration, and intolerance to EN feeding.

AUTHOR DISCLOSURES

The authors declare that they have no conflicts of interest.

This study was supported by grants from the open project of national clinical research center for geriatric diseases (NCRCP-PLAGH-2017008), the medical big data research project of PLA general hospital (2016MBD-014) and the class general financial grant from the china postdoctoral science foundation (2016M592985).

REFERENCES

- Bouffard YH, Delafosse BX, Annat GJ, Viale JP, Bertrand OM, Motin JP. Energy expenditure during severe acute pancreatitis. *JPEN J Parenter Enteral Nutr.* 1989;13:26-9. doi: 10.1177/014860718901300126.
- Li Q, Zhang Z, Xie B, Ji X, Lu J, Jiang R et al. Effectiveness of enteral feeding protocol on clinical outcomes in critically ill patients: a before and after study. *PLoS One.* 2017;12:e0182393. doi: 10.1371/journal.pone.0182393.
- Wikjord K, Dahl V, Sovik S. Effects on nutritional care practice after implementation of a flow chart-based nutrition support protocol in an intensive care unit. *Nursing Open.* 2017;4:282-91. doi:10.1002/nop2.99.
- Kim SH, Park CM, Seo JM, Choi M, Lee DS, Chang DK et al. The impact of implementation of an enteral feeding protocol on the improvement of enteral nutrition in critically ill adults. *Asia Pac J Clin Nutr.* 2017;26:27-35. doi: 10.6133/apjcn.122015.01.
- Barr J, Hecht M, Flavin KE, Khorana A, Gould MK. Outcomes in critically ill patients before and after the implementation of an evidence-based nutritional management protocol. *Chest.* 2004;125:1446-57.
- Kuslapuu M, Jogela K, Starkopf J, Reintam Blaser A. The reasons for insufficient enteral feeding in an intensive care unit: A prospective observational study. *Intensive & critical care nursing.* 2015;31:309-14. doi: 10.1016/j.iccn.2015.03.001.
- Taylor BE, McClave SA, Martindale RG, Warren MM, Johnson DR, Braunschweig C et al. Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *Crit Care Med.* 2016;44:390-438. doi: 10.1097/ccm.0000000000001525.
- Kreymann KG, Berger MM, Deutz NE, Hiesmayr M, Jolliet P, Kazandjiev G et al. ESPEN Guidelines on Enteral Nutrition: Intensive care. *Clin Nutr.* 2006;25:210-23. doi: 10.1016/j.clnu.2006.01.021.
- Patel JJ, Lemieux M, McClave SA, Martindale RG, Hurt RT, Heyland DK. Critical care nutrition support best practices: key differences between Canadian and American Guidelines. *Nutr Clin Pract.* 2017;32:633-44. doi: 10.1177/0884533617722165.
- Hales BM, Pronovost PJ. The checklist--a tool for error management and performance improvement. *J Crit Care.* 2006;21:231-5. doi: 10.1016/j.jcrc.2006.06.002.
- Winters BD, Gurses AP, Lehmann H, Sexton JB, Rampersad CJ, Pronovost PJ. Clinical review: checklists - translating evidence into practice. *Crit Care.* 2009;13:210. doi: 10.1186/cc7792.
- Brunsveld-Reinders AH, Arbous MS, Kuiper SG, de Jonge E. A comprehensive method to develop a checklist to increase safety of intra-hospital transport of critically ill patients. *Crit Care.* 2015;19:214. doi: 10.1186/s13054-015-0938-1.
- Taylor JE, McDonald SJ, Earnest A, Buttery J, Fusinato B, Hovenden S, Wallace A, Tan K. A quality improvement initiative to reduce central line infection in neonates using checklists. *Eur J Pediatr.* 2017;176:639-46. doi: 10.1007/s00431-017-2888-x.
- Pamplin J, Kuwamoto R, Bradstreet H, Linfoot J, Chung K, Grathwohl K. A daily checklist can change intensive care unit hand hygiene culture. *Crit Care Med.* 2010;38:U164.
- Howie WO, Dutton RP. Implementation of an evidence-based extubation checklist to reduce extubation failure in patients with trauma: a pilot study. *AANA Journal.* 2012;80:179-84.
- Ba RRW. Using aspiration risk score in department of geriatrics. *Today Nurse.* 2017;2:142-4.
- McCullough JPA, Lipman J, Presneill JJ. The statistical curriculum within randomized controlled trials in critical illness. *Crit Care Med.* 2018;46:1985-90. doi: 10.1097/ccm.0000000000003380.
- Orinovskiy I, Raizman E. Improvement of nutritional intake in intensive care unit patients via a nurse-led enteral nutrition feeding protocol. *Crit Care Nurse.* 2018;38:38-44. doi: 10.4037/ccn2018433.
- Blaser AR, Starkopf J, Kirsimagi U, Deane AM. Definition, prevalence, and outcome of feeding intolerance in intensive care: a systematic review and meta-analysis. *Acta Anaesthesiol Scand.* 2014;58:914-922. doi: 10.1111/aas.12302.
- Mancl EE, Muzevich KM. Tolerability and safety of enteral nutrition in critically ill patients receiving intravenous vasopressor therapy. *JPEN J Parenter Enteral Nutr.* 2013; 37:641-51. doi: 10.1177/0148607112470460.
- Hu B, Sun R, Wu A, Ni Y, Liu J, Guo F et al. Severity of acute gastrointestinal injury grade is a predictor of all-cause mortality in critically ill patients: a multicenter, prospective, observational study. *Crit Care.* 2017;21:188. doi: 10.1186/s13054-017-1780-4.
- Li H, Zhang D, Wang Y, Zhao S. Association between acute gastrointestinal injury grading system and disease severity and prognosis in critically ill patients: A multicenter, prospective, observational study in China. *J Crit Care.* 2016; 36:24-8. doi: 10.1016/j.jcrc.2016.05.001.
- Dhaliwal R, Cahill N, Lemieux M, Heyland DK. The Canadian critical care nutrition guidelines in 2013: an update on current recommendations and implementation

- strategies. *Nutr Clin Pract.* 2014;29:29-43. doi: 10.1177/0884533613510948.
24. Haskins IN, Baginsky M, Gamsky N, Sedghi K, Yi S, Amdur RL, Gergely M, Sarani B. Volume-based enteral nutrition support regimen improves caloric delivery but may not affect clinical outcomes in critically ill patients. *JPEN J Parenter Enteral Nutr.* 2017;41:607-11. doi: 10.1177/0148607115617441.
25. Weiss CH, Moazed F, McEvoy CA, Singer BD, Szeleifer I, Amaral LAN et al. Prompting physicians to address a daily checklist and process of care and clinical outcomes a single-site study. *Am J Respir Crit Care Med.* 2011;184:680-6. doi: 10.1164/rccm.201101-0037OC.
26. Centofanti JE, Duan EH, Hoad NC, Swinton ME, Perri D, Waugh L, Cook DJ. Use of a daily goals checklist for morning ICU rounds: a mixed-methods study. *Crit Care Med.* 2014;42:1797-803. doi: 10.1097/ccm.0000000000000331.

Supplemental table 1. Aspiration risk score

Scores	1	2	3
Age	10-49 years	50-80 years	>80 years or <10 years
Consciousness	Conscious	Conscious and sedation	Coma
Sputum	Little	More and thickness	More and thin
Alzheimer's disease, cerebrovascular accident, myasthenia gravis, Parkinson's disease	None	One	More than one
Diet	Abrosia	Normal	Liquid or semiliquid diets
Body position	Semireclining position $\geq 30^\circ$	Semireclining position $< 30^\circ$	Horizontal position
Water swallow test	1 grade	2 grade	≥ 3 grade
Artificial airway and mechanical ventilation	None	Positive	/
Aspiration history	None	/	Positive

Aspiration risk are classified as: Low, 0-10; High, 11-26