

Original Article

Citrus fruit intake and the risk of nasopharyngeal carcinoma

Xin-Xin Feng MSc^{1,2,3}, Ming-Xu Wang MD³, Ming Li MD⁴, Xiao Tang MSc³, Hong Jiang MD³, Rui Wang MSc³, Le Ma MD³, Yue Yin MD³, Chang-Rui Wu MD¹

¹The First Affiliated Hospital of Xi'an Jiaotong University Health Science Center, Xi'an, China

²Key Laboratory of Shaanxi Province for Craniofacial Precision Medicine Research, Hospital of Stomatology, Xi'an Jiaotong University, Xi'an, China

³School of Public Health, Xi'an Jiaotong University Health Science Center, Xi'an, China

⁴Center for Population Health Research, Division of Health Sciences, University of South Australia, Adelaide, Australia

Background and Objectives: Citrus fruit are suggested to be associated with reduced risk of nasopharyngeal carcinoma (NPC), but findings from epidemiologic studies have been inconsistent. We aimed to synthesize the association by conducting a meta-analysis of existing evidence. **Methods and Study Design:** Databases including Medline, EMBASE, Web of science, and the Cochrane Library were searched for eligible studies up to March 2019 using a series comprehensive searching terms. The adjusted odds ratios (ORs) of citrus fruit intake with NPC risk from each study were extracted to calculate a pooled association estimate with its 95% confidence interval (CI). **Results:** Nine studies totaling 3304 cases and 3850 controls were included in this analysis. Citrus fruit intake was significantly associated with reduced risk of NPC (OR: 0.72, 95% CI 0.58-0.91, $p=0.005$). In addition, this association tended to be stronger in Chinese (OR: 0.67, 95% CI 0.54-0.84, $p<0.001$). Dose-response analysis using cubic splines showed the risk of NPC decreased by 21% for citrus fruit intake of 4 times/week (OR: 0.79, 95% CI 0.66-0.94). **Conclusions:** The consumption of citrus fruit was associated with a significant reduced risk of NPC, especially in Chinese population.

Key Words: citrus fruit consumption, nasopharyngeal carcinoma, dose-response analysis, protective effects, meta-analysis

INTRODUCTION

Nasopharyngeal carcinoma (NPC) is a squamous-cell carcinoma that originates in the epithelial lining of the nasopharynx.¹ It is more prevalent in populations in Southeast Asia, the Arctic, and the Middle East/North Africa,^{2,3} and more than 70% of new reported cases were from eastern/south-east Asia in 2012.⁴ The overall age-standardized incidence rate and mortality of NPC in 2013 were 25/100,000 and 14/100,000 amongst people living in South China, higher than Caucasians from America and other western countries with the incidence rate and mortality of 1/100,000.^{2,5} To date, the exact pathogenic mechanism of NPC has not been understood. However, the distinctive ethnic distribution of the carcinoma worldwide suggests that besides genetic susceptibility and Epstein-Barr virus (EBV) infection, environmental factors might be associated with the risk of NPC.^{6,7}

A number of factors have been suggested to be associated with increased risk of NPC, including the consumption of Cantonese-style salted fish,^{8,9} preserved vegetables,^{10,11} cigarette smoking,¹² and alcohol drinking.¹³ Whereas, fruit consumption has been associated with a reduced risk of NPC.¹⁴ Citrus fruit contains a variety of antioxidants such as vitamin C, folate, and carotenoids,

and other bioactive components, which are believed to have protective effects on cancer that attributed to antioxidant activities, prevention of nitrosamine formation, and other anticarcinogenic properties.¹⁵

Previous studies have demonstrated that citrus fruit consumption are associated with decreased risk of various cancers from gastrointestinal and urogenital systems.^{16,17} Over the decades, there have been studies reported controversial association between citrus fruit consumption and the risk of NPC.¹⁸⁻²⁶ In addition, variations in geographic location, methodology, and adjustment factors across studies may have differential impacts on NPC risk, yet the directions of these impacts have not been investigated.

We conducted a meta-analysis of all published studies to evaluate the association between citrus fruit intake and

Corresponding Author: Dr Chang-Rui Wu, The First Affiliated Hospital of Xi'an Jiaotong University Health Science Center, 277 Yanta West Road, Xi'an, Shaanxi, 710061, China.

Tel: +86-15802994257; Fax: +86-29-82655032

Email: gxwcr001@163.com

Manuscript received 05 May 2019. Initial review completed 04 June 2019. Revision accepted 28 July 2019.

doi:

NPC risk. We also quantified dose-response relations of citrus fruit intake with NPC risk.

METHODS

The study followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines²⁷ and the Meta-analysis of Observational Studies in Epidemiology (MOOSE) group checklist.²⁸

Search strategy

Databases of Medline, EMBASE, Web of science, and the Cochrane Library were searched for published studies till March 2019. The following search terms were applied to maximize sensitivity: 'diet' OR 'fruit' OR 'citrus' OR 'orange' OR 'tangerine' OR 'grapefruit' OR 'lemon' OR 'lime' and 'nasopharyngeal neoplasms' OR 'nasopharyngeal carcinoma' OR 'nasopharyngeal cancer' OR 'nasopharynx cancer' OR 'pharyngeal cancer' OR 'NPC'. We also manually searched the references listed in the retrieved articles and published systematic reviews to identify additional relevant studies. Potential unpublished studies were searched by contacting corresponding authors of relevant primary studies.

Study selection

An initial screen of titles and abstracts of the identified studies was conducted for eligibility. Full text of the relevant publications were retrieved based on the following study eligibility criteria: (1) the primary outcome was NPC; (2) the exposure was citrus fruit consumption (including oranges, tangerines, grapefruits, lemons and limes, administered alone or in combination); (3) odds ratios (ORs) and its 95% confidence intervals (CIs) for the association between citrus fruit intake and NPC risk (or sufficient data to calculate them) was reported; and (4) potential confounders were controlled by matching or multivariable analysis in the studies. If data were duplicated in more than one study, only the most updated data were chosen for analysis. Two investigators (X.-X.F. and X.T.) assessed the studies independently, and any discrepancies were resolved by group discussion among all co-authors until a consensus was reached.

Data extraction and quality assessment

Data were extracted independently by two investigators (X.-X.F. and X.T.) using a standardized form. The data included: study characteristics (first author, publication year, country, study design), participant characteristics (the number of cases/subjects, sex distribution, age range or mean age), dietary assessment methods, comparison of exposure level and their effects estimates, diagnosis method of NPC, and variables adjusted in the analysis. If a study provided several risk estimates, the maximum adjusted estimate was used. When ORs were not available, crude effect estimates and 95% CIs were calculated using the primary data. As one study did not report any association for the highest category of citrus fruit intake, we used the OR from the adjacent category.²³

The quality of the studies was assessed using the Newcastle-Ottawa scale (NOS) based on the selection of study groups (four criteria), the comparability of study groups (one criterion), and the assessment of the exposure (three

criteria).²⁹ Each study was assigned 0-9 points. A study was regarded high in quality if it was scored 6 or more, and those scored less than 6 were considered low quality.³⁰ Any discrepancies in data extraction and quality assessment were resolved through group discussion.

Statistical analysis

Pooled ORs with the corresponding 95% CIs were calculated to measure the strength of the association between citrus fruit intake and the risk of NPC. Heterogeneity across the studies was evaluated by the DerSimonian and Laird's Q statistic ($p < 0.10$ indicated statistically significant heterogeneity) and I² statistic (values of 25%, 50% and 75% were defined to represent low, moderate and high heterogeneity, respectively).^{31,32} Random-effects model was used to calculate the pooled ORs when substantial heterogeneity was detected; while the fixed-effects model was adopted otherwise.³³ Forest plot was generated to visualize the ORs and the 95% CIs across studies. The source of heterogeneity was explored with stratification and meta-regression analyses by geographic location (China vs not China), sources of controls (population-based vs hospital-based), diagnosis method (pathology reports vs the clinicians), smoking status (adjusted vs not adjusted), alcohol drinking (adjusted vs not adjusted), and quality score (high vs low).

In addition, we assessed the potential dose-response relation between citrus fruit intake and the risk of NPC using the method proposed by Greenland and Longnecker.³⁴ Within each study, the median or mean citrus fruit intake in each category was assigned to the corresponding OR estimate. The midpoint of the upper and lower bounds in each citrus fruit category was utilized if median or mean was not reported in the study. The open-ended categories were assumed to have the same amplitude as the adjacent categories. A restricted cubic spline model with 3 fixed knots at 10%, 50%, and 90% through the total distribution of the reported intake and the generalized least-squares regression were used to estimate the nonlinear association.³⁵

Sensitivity analyses were performed by removing one study at a time to examine whether the overall result was affected significantly by any single study. Publication bias was evaluated by using visual inspection of a funnel plot, and additionally evaluated statistically with Egger and Begg's tests.^{36,37}

All analyses were conducted by using Stata version 10.0 (Stata Corporation, College Station, TX, USA). All *p* values were 2-tailed with a statistically significance level of 0.05.

RESULTS

Search results

A total of 2558 citations were obtained from the initial search, of which 904 were from Medline, 640 from EMBASE, 989 from Web of Science, and 25 from the Cochrane Library. Fifty-five full texts were potentially eligible after removing duplicates and screening titles or abstracts. Of these, nine studies met the selection criteria and were included in the present meta-analysis (Figure 1; Supplemental table 1).¹⁸⁻²⁶




Figure 1. The flow diagram of study selection from MEDLINE (www.ncbi.nlm.nih.gov/pubmed), EMBASE (<http://www.embase.com/>), Web of Science (<http://thomsonreuters.com/web-of-science/>), and the Cochrane Library (<http://www.cochranelibrary.com/>).

Study characteristics

The main characteristics of included studies are summarized in Table 1. Of the nine studies, five were conducted in China, two in Europe, one in the United States, and one in North Africa. The study samples ranged from 135 to 1967, giving a total of 7154 participants composed of 3304 NPC cases. All studies consisted of both sexes and the proportion of males varied between 50.0% and 79.3%. As to controls, five studies were population-based,^{18-21,25} and the others were hospital-based.^{22-24,26} Citrus fruit intake was assessed by food frequency questionnaires (FFQ) in all studies via self-administrated survey¹⁹ or face-to-face interviews.^{18,20-26} NPC was ascertained based on pathological diagnosis in eight studies,^{18-21,23-26} or on clinical diagnosis in one.²² All of the studies were matched or adjusted for age, and other factors included sex,^{18,20-26} smoking status,^{19,21,24-26} education,^{21,24-26} exposure to potential toxic substances,^{21,22,24,25} energy intake,^{19,24-26} residence,^{20,23,26} alcohol drinking,^{19,26} chronic ear and nose condition,^{21,24} and NPC family history.²⁵ Overall, eight studies out of nine were of high quality, with a score of ≥ 6 .

Citrus fruit intake and NPC

Among the nine studies, four reported a significantly decreased risk of NPC associated with citrus fruit in-

take,^{18,20,21,24} and the comparison of exposure level and their effects estimates in each primary study were shown in Supplemental Table 2. Significant heterogeneity was observed across the studies ($I^2=51.6\%$; $p_{\text{heterogeneity}}=0.04$). Pooled OR was 0.72 (95% CI 0.58-0.91, $p=0.005$) for the highest relative to the lowest citrus fruit intake using the random-effects model (Figure 2). In the dose-response analysis using cubic splines, the risk of NPC decreased by 21% with increasing intake of citrus fruit up to 4 times/week (OR: 0.79, 95% CI 0.66-0.94, Figure 3). No benefit was apparent when increasing intake above this value.

When stratified by geographic location, studies conducted in China tended to report a stronger association of citrus fruit intake with NPC risk (OR: 0.67, 95% CI 0.54-0.84, $p<0.001$), whereas the association was nonsignificant in other populations (OR: 0.84, 95% CI 0.50-1.42, $p=0.515$). Subgroup analysis across other study and participant characteristics found these factors did not significantly alter the shape of the association (Table 2).

Sensitivity analysis and publication bias

Sensitivity analyses showed that the pooled ORs for NPC ranged from 0.67 (95% CI 0.57-0.80) when the study conducted by Feng et al²² was excluded to 0.76 (95% CI 0.59-0.97) when the study conducted by Yuan et al.²¹ was excluded, which indicate that the results of the present meta-analysis remained stable (Supplemental Table 3). No publication bias was suggested, either by the funnel plot or by statistical tests (Begg's test, $p=0.60$; and Egger's test, $p=0.91$; Supplemental figure 1).

DISCUSSION

The findings from the present meta-analysis suggested that high intake of citrus fruit was associated with a significant decreased risk of NPC. In addition, a significant dose-responsive relation was observed, which indicating a reduction risk of 21% NPC for citrus fruit intake of 4 times/week. The results also showed that studies conducted in China tended to report a stronger association.

The association between citrus fruit intake and cancers have been examined. For example, Wang et al. noted that a decreased risk of esophageal cancer was associated with higher intake of citrus fruit (OR: 0.63, 95% CI 0.52-0.75) in a meta-analysis of 6 cohort studies and 13 case-control studies.³⁸ Similarly, a decreased risk of 50% was confirmed by Bravi et al for oral and pharyngeal cancer among those with highest quintile intakes of citrus fruit after adjustment of age, sex, center (Italy, Switzerland), education, year of interview, BMI, tobacco smoking, alcohol drinking and energy intake.³⁹ Citrus fruit has been shown to be protective as well for pancreatic cancer and breast cancer.^{40,41} The present results showed that higher consumption of citrus fruit was associated with a reduced risk of NPC, adding in the evidence that citrus fruit intake would have a protective effect on cancer development.

Our finding of a decreased NPC risk with high citrus fruit intake is biologically plausible, and the potential mechanisms are consistent with numerous proposed mechanisms whereby citrus fruit decrease risk for cancers because of the similar pathogenesis. Citrus fruits are rich

Table 1. Characteristics of studies on citrus fruit intake and NPC risk

First author, year	Country	Study design	Case/controls (n)	Sex (%male)	Age (years)	Dietary assessment	Diagnosis method	Adjustments	Quality score [†]
Chen et al, 1997 ¹⁸	China	Case-control (PCC)	104/104	50.0	≤55	FFQ-33 face-to-face interview	Pathology reports	Age, sex	5
Farrow et al, 1998 ¹⁹	USA	Case-control (PCC)	133/212	67.5	18-74	FFQ-100 self-administrated survey	Pathology reports	Age, alcohol consumption, cigarette smoking, total caloric intake	6
Armstrong et al, 1998 ²⁰	China	Case-control (PCC)	282/282	69.1	45 (mean)	FFQ-55 face-to-face interview	Pathology reports	Age, sex, residence, marital status	7
Yuan et al, 2000 ²¹	China	Case-control (PCC)	935/1032	69.5	15-74	FFQ-38 face-to-face interview	Pathology reports	Age, gender, level of education, cigarette smoking, exposure to smoke from heated rapeseed oil and burning coal during cooking, occupational exposure to chemical fumes, history of chronic ear and nose condition	7
Feng et al, 2007 ²²	North Africa	Case-control (HCC)	636/615	69.5	15-81	FFQ face-to-face interview	The clinicians	Sex, center, age, socio-economic status variables, exposure to toxic substances	8
Nešić1 et al, 2010 ²³	Serbia	Case-control (HCC)	45/90	64.4	NR	FFQ-100 face-to-face interview	Pathology reports	Age, sex, residence	6
Liu et al, 2012 ²⁴	China	Case-control (HCC)	600/600	74.7	47 (mean)	FFQ-78 face-to-face interview	Pathology reviews	Age, gender, BMI, educational level, marital status, occupation, household income, occupational, domestic exposure to potential toxic substances, chronic rhinitis history, smoking status, passive smoking, daily energy intake, energy-adjusted intakes of other food groups	7
Hsu et al, 2012 ²⁵	China	Case-control (PCC)	371/321	69.4	≤75	FFQ-66 face-to-face interview	Pathology reports	Age, gender, ethnicity, educational level, NPC family history, total calories intake, years of cigarette smoking, exposures to formaldehyde and wood dust	7
Polesel et al, 2013 ²⁶	Italy	Case-control (HCC)	198/594	79.3	18-76	FFQ-78 face-to-face interview	Pathology reports	Center, sex, age, place of living, year of interview, education, tobacco smoking, alcohol drinking, non-alcohol energy	7

BMI: body Mass Index; FFQ: food frequency questionnaires; HCC: hospital-based case-control study; NPC: nasopharyngeal carcinoma; NR: not reported; PCC: population-based case-control study.

[†]Study quality was assessed based on the validated Newcastle-Ottawa scale; studies with ≥6 were considered high-quality and low otherwise.³⁰

Table 2. Stratified analysis of the association between citrus fruit intake and NPC[†]

Subgroup	n	Case/controls (n)	Pooled OR	95% CI	P	
					p_h^{\dagger}	p_h^{\ddagger}
Geographic location						
China	5	2292/2339	0.67	0.54, 0.84	0.13	0.32
Not China	4	1012/1511	0.84	0.50, 1.42	0.09	
Sources of controls						
Population-based	5	1825/1951	0.67	0.53, 0.86	0.13	0.46
Hospital-based	4	1479/1899	0.81	0.50, 1.32	0.04	
Diagnosis method						
Pathology reports	8	2668/3235	0.67	0.57, 0.80	0.30	NP
The clinicians	1	636/615	1.50	0.90, 2.60	NA	
Smoking status						
Adjusted	5	2237/2759	0.72	0.56, 0.91	0.21	0.94
Not adjusted	4	1067/1091	0.72	0.43, 1.21	0.01	
Alcohol drinking						
Adjusted	2	331/806	0.73	0.48, 1.10	0.83	>0.99
Not adjusted	7	2973/3044	0.72	0.55, 0.96	0.01	
Quality score [§]						
High	8	3200/3746	0.74	0.56, 0.97	0.02	NP
Low	1	104/104	0.66	0.50, 0.89	NA	

CI: confidence interval; NA: not applicable (because only one study); NP: meta-regression was not possible; NPC: nasopharyngeal carcinoma; OR: odds ratio.

[†] p_h values were for heterogeneity within a subgroup.

[‡] p_h values were for heterogeneity between subgroups by meta-regression.

[§]Study quality was assessed based on the validated Newcastle-Ottawa scale; studies with ≥ 6 were considered high-quality and low otherwise.³⁰




Figure 2. Forest plot on the association between citrus fruit intake (highest-versus-lowest category) and nasopharyngeal carcinoma (NPC). The boxes and lines indicate the odds ratios (ORs) and their confidence intervals (CIs) on a log scale for each study. The pooled odds ratio (OR) is represented by a diamond. The size of the black squares indicates the relative weight of each estimate. The arrow indicates that the lowest confidence interval of the OR is below the range of the axis.

in multiple cancer chemopreventive agents, including antioxidants, limonoids and furocoumarins. As principal antioxidants, vitamin C may play a role in the prevention of different malignant diseases development including NPC by blocking the metabolic activation of carcinogens, stimulating the immune function, and preventing oxidative stress.^{42,43} Besides decreasing oxidative damage to DNA, carotenoids and folate are implicated in the etiology of cancer via effects on inhibition of cell proliferation and differentiation, repair and methylation of DNA and

scavenging free radicals.^{44,45} Moreover, as prominent phytochemicals in citrus fruit, limonoids appear to possess substantial antineoplastic activity against chemically induced cancers,⁴⁶ as animal experiment has demonstrated that limonoids were positively correlated with the induction of Glutathione S-transferase (GST) activity to catalyzes the conjugation of glutathione with electrophilic xenobiotics that induce activated carcinogens.⁴⁷ Notably, furanocoumarins had been indicated to exhibit several antiproliferative activities against the growth of cancer




Figure 3. Odds ratios (ORs) and the corresponding 95% confidence intervals (CIs) for the dose–response relationship between citrus fruit intake and nasopharyngeal carcinoma (NPC) risk. The circles represent ORs of varied doses in each primary study. The solid line represents best-fitting cubic spline model and the dotted lines represent 95% CIs.

cells through modulation of several molecular pathways, such as upregulation of the signal transducer and activator of transcription 3, phosphatidylinositol-3-kinase/protein kinase B, nuclear factor- κ B, and mitogen-activated protein kinase expression.⁴⁸

It should be noted that citrus fruit appeared to have protective trend on NPC in Chinese populations. Familial NPC, which is common in China, was linked to higher susceptibility to genetic locus mutation,⁴⁹ which warrants further investigation, as most of the studies ascertained in this meta-analysis did not provide information regarding the family history of nasopharyngeal cancer. Higher intakes of fruit has been show to attenuate the genetic predisposition to obesity,⁵⁰ which is associated with increased risk of NPC in a prospective cohort study from Israel and two case-control studies from Zhejiang and Guangdong, China.⁵¹ On the other hand, preserved foods such as Cantonese salted fish, preserved vegetables and eggs, which are consumed frequently in south China, contain higher nitrosamines that were postulated as carcinogenic in the development of NPC over time.⁵² Thus, the protective effect of citrus fruit on NPC risk via prevention of nitrosamine formation is more likely to be detected among Chinese. Finally, citrus extracts has been indicated to have inhibitory effects on EBV early antigen activation induced by the tumor promoter in an in vitro experiment.⁵³ The epidemiology of childhood EBV infection is different in Chinese and other populations, and most children in the developing countries are infected with EBV before the age of three.⁵⁴ Therefore, the effect of citrus fruit could be different depending on the timing of EBV infection.

The limitations of this meta-analysis should be noted. First, the present study was based on case-control studies and are susceptible to the inherent recall bias and selection bias, which may overstate the association.⁵⁵ However, the quantitative analyses in our study were based on a large number of participants and most studies were evaluated to be high quality using the NOS,²⁹ which suggested the results were quite robust. Second, the associations across studies are heterogeneous, indicating differences in study regions, source of controls, and adjusted confound-

ers; however, stratified analyses by these factors did not affect the association of citrus fruit intake with NPC risk in the present meta-analysis. Third, our inability to standardize citrus fruit intake in all of the included studies might have biased the results, although the likelihood should be small because the reference group and the highest citrus fruit intake group were similar in most primary studies. Fourth, residual or unknown confounders could not be completely excluded. For example, higher fruit intakes is more likely to related to certain healthy behaviors that would bias the true associations.⁵⁶ Fifth, possible measurement errors in the dietary assessment using FFQ that represent a subjective approximation of past dietary behaviors rather than an assessment of absolute intakes in the included studies may affect the effect estimates, but the directions of these impacts are unpredictable. Sixth, although no publication bias was suggested visually or statistically, it cannot be fully ruled out.

Conclusion

In summary, this meta-analysis suggested that citrus fruit intake was significantly associated with a lower risk of NPC, particularly in Chinese populations. Further well-designed cohort studies and randomized controlled trials including other populations are warranted to confirm the impact of citrus fruit on NPC risks.

AUTHOR DISCLOSURES

The authors declare no conflict of interest.


This study was supported by the Key Laboratory of Shaanxi Province for Craniofacial Precision Medicine Research, College of Stomatology, Xi'an Jiaotong University (2016LHM-KFKT009), the New-star Plan of Science and Technology of Shaanxi Province (2015LJXX-07), the China Postdoctoral Science Special Foundation (2015T81036), the Fundamental Research Funds for the Central Universities (qngz2016004), the Natural Science Foundation of Shaanxi Province of China (2017JM8041), and the Fundamental Research Funds for the Central Universities (zdyf2017007).

REFERENCES

1. Wei WI, Sham JS. Nasopharyngeal carcinoma. *Lancet*. 2005; 365:2041-54. doi: 10.1016/s0140-6736(05)66698-6.

2. Tang LL, Chen WQ, Xue WQ, He YQ, Zheng RS, Zeng YX, Jia WH. Global trends in incidence and mortality of nasopharyngeal carcinoma. *Cancer Lett.* 2016;374:22-30. doi: 10.1016/j.canlet.2016.01.040.
3. Chang ET, Adami HO. The enigmatic epidemiology of nasopharyngeal carcinoma. *Cancer Epidemiol Biomarkers Prev.* 2006;15:1765-77. doi: 10.1158/1055-9965.epi-06-0353.
4. Shield KD, Ferlay J, Jemal A, Sankaranarayanan R, Chaturvedi AK, Bray F, Soerjomataram I. The global incidence of lip, oral cavity, and pharyngeal cancers by subsite in 2012. *CA Cancer J Clin.* 2017;67:51-64. doi: 10.3322/caac.21384.
5. Wei KR, Zheng RS, Zhang SW, Liang ZH, Li ZM, Chen WQ. Nasopharyngeal carcinoma incidence and mortality in China, 2013. *Chin J Cancer.* 2017;36:90. doi: 10.1186/s40880-017-0257-9.
6. Lo KW, To KF, Huang DP. Focus on nasopharyngeal carcinoma. *Cancer Cell.* 2004;5:423-8. doi: 10.1016/S1535-6108(04)00119-9.
7. Bei JX, Li Y, Jia WH, Feng BJ, Zhou G, Chen LZ, Feng QS, Low HQ, Zhang H, He F, Tai ES, Kang T, Liu ET, Liu J, Zeng YX. A genome-wide association study of nasopharyngeal carcinoma identifies three new susceptibility loci. *Nat Genet.* 2010;42:599-603. doi: 10.1038/ng.601.
8. Yu MC, Ho JH, Lai SH, Henderson BE. Cantonese-style salted fish as a cause of nasopharyngeal carcinoma: report of a case-control study in Hong Kong. *Cancer Res.* 1986;46:956-61. doi: 10.1016/0304-3835(86)90091-1.
9. Sriamporn S, Vatanasapt V, Pisani P, Yongchaiyudha S, Rungpitarangsi V. Environmental risk factors for nasopharyngeal carcinoma: a case-control study in northeastern Thailand. *Cancer Epidemiol Biomarkers Prev.* 1992;1:345-8. doi: 10.1080/00071669208417493.
10. Lee HP, Gourley L, Duffy SW, Esteve J, Lee J, Day NE. Preserved foods and nasopharyngeal carcinoma: a case-control study among Singapore Chinese. *Int J Cancer.* 1994;59:585-90. doi: 10.1002/ijc.2910590502.
11. Gallicchio L, Matanoski G, Tao XG, Chen L, Lam TK, Boyd K, Robinson KA, Balick L, Mickelson S, Caulfield LE, Herman JG, Guallar E, Alberg AJ. Adulthood consumption of preserved and nonpreserved vegetables and the risk of nasopharyngeal carcinoma: a systematic review. *Int J Cancer.* 2006;119:1125-35. doi: 10.1002/ijc.21946.
12. Chow WH, McLaughlin JK, Hrubec Z, Nam JM, Blot WJ. Tobacco use and nasopharyngeal carcinoma in a cohort of US veterans. *Int J Cancer.* 1993;55:538-40. doi: 10.1002/ijc.2910550403.
13. Li Y, Yang H, Cao J. Association between alcohol consumption and cancers in the Chinese population—a systematic review and meta-analysis. *PLoS One.* 2011;6:e18776. doi: 10.1371/journal.pone.0018776.
14. Yu MC, Huang TB, Henderson BE. Diet and nasopharyngeal carcinoma: a case-control study in Guangzhou, China. *Int J Cancer.* 1989;43:1077-82. doi: 10.1002/ijc.2910430621.
15. Silalahi J. Anticancer and health protective properties of citrus fruit components. *Asia Pac J Clin Nutr.* 2002;11:79-84. doi: 10.1046/j.1440-6047.2002.00271.x.
16. Bae JM, Kim EH. Dietary intakes of citrus fruit and risk of gastric cancer incidence: an adaptive meta-analysis of cohort studies. *Epidemiol Health.* 2016;38:e2016034. doi: 10.4178/epih.e2016034.
17. Liang S, Lv G, Chen W, Jiang J, Wang J. Citrus fruit intake and bladder cancer risk: a meta-analysis of observational studies. *Int J Food Sci Nutr.* 2014;65:893-8. doi: 10.3109/09637486.2014.917151.
18. Chen DL, Huang TB. A case-control study of risk factors of nasopharyngeal carcinoma. *Cancer Lett.* 1997;117:17-22. doi: 10.1016/S0304-3835(97)00182-1.
19. Farrow DC, Vaughan TL, Berwick M, Lynch CF, Swanson GM, Lyon JL. Diet and nasopharyngeal cancer in a low-risk population. *Int J Cancer.* 1998;78:675-9. doi: 10.1002/(sici)1097-0215(19981209)78:6<675::aid-ijc2>3.0.co;2-j.
20. Armstrong RW, Imrey PB, Lye MS, Armstrong MJ, Yu MC, Sani S. Nasopharyngeal carcinoma in Malaysian Chinese: salted fish and other dietary exposures. *Int J Cancer.* 1998;77:228-35. doi: 10.1002/(sici)1097-0215(19980717)77:2<228::aid-ijc11>3.0.co;2-7.
21. Yuan JM, Wang XL, Xiang YB, Gao YT, Ross RK, Yu MC. Preserved foods in relation to risk of nasopharyngeal carcinoma in Shanghai, China. *Int J Cancer.* 2000;85:358-63. doi: 10.1002/(sici)1097-0215(20000201)85:3<358::aid-ijc11>3.0.co;2-e.
22. Feng BJ, Jalbout M, Ayoub WB, Khyatti M, Dahmoul S, Ayad M et al. Dietary risk factors for nasopharyngeal carcinoma in Maghreb countries. *Int J Cancer.* 2007;121:1550-5. doi: 10.1002/ijc.22813.
23. Nesic V, Sipetic S, Vlajinac H, Stosic-Divjak S, Jesic S. Risk factors for the occurrence of undifferentiated carcinoma of nasopharyngeal type: a case-control study. *Srp Arh Celok Lek.* 2010;138:6-10. doi: 10.2298/SARH1002006N.
24. Liu YT, Dai JJ, Xu CH, Lu YK, Fan YY, Zhang XL, Zhang CX, Chen YM. Greater intake of fruit and vegetables is associated with lower risk of nasopharyngeal carcinoma in Chinese adults: a case-control study. *Cancer Causes Control.* 2012;23:589-99. doi: 10.1007/s10552-012-9923-z.
25. Hsu WL, Pan WH, Chien YC, Yu KJ, Cheng YJ, Chen JY et al. Lowered risk of nasopharyngeal carcinoma and intake of plant vitamin, fresh fish, green tea and coffee: a case-control study in Taiwan. *PLoS One.* 2012;7:e41779. doi: 10.1371/journal.pone.0041779.
26. Polesel J, Serraino D, Negri E, Barzan L, Vaccher E, Montella M et al. Consumption of fruit, vegetables, and other food groups and the risk of nasopharyngeal carcinoma. *Cancer Causes Control.* 2013;24:1157-65. doi: 10.1007/s10552-013-0195-z.
27. Moher D, Liberati A, Tetzlaff J, Altman DG, Grp P. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *Annals of Internal Medicine.* 2009;151:264-W64. doi: 10.7326/0003-4819-151-4-200908180-00135.
28. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, Moher D, Becker BJ, Sipe TA, Thacker SB. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA.* 2000;283:2008-12. doi: 10.1001/jama.283.15.2008.
29. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol.* 2010;25:603-5. doi: 10.1007/s10654-010-9491-z.
30. Li F, Duan F, Zhao X, Song C, Cui S, Dai L. Red meat and processed meat consumption and nasopharyngeal carcinoma risk: a dose-response meta-analysis of observational studies. *Nutr Cancer.* 2016;68:1034-43. doi: 10.1080/01635581.2016.1192200.
31. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med.* 2002;21:1539-58. doi: 10.1002/sim.1186.

32. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ*. 2003;327:557-60. doi: 10.1136/bmj.327.7414.557.
33. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986;7:177-88. doi: 10.1016/0197-2456(86)90046-2.
34. Greenland S, Longnecker MP. Methods for trend estimation from summarized dose-response data, with applications to meta-analysis. *Am J Epidemiol*. 1992;135:1301-9. doi: 10.1093/oxfordjournals.aje.a116237.
35. Durrleman S, Simon R. Flexible regression models with cubic splines. *Stat Med*. 1989;8:551-61. doi: 10.1002/sim.4780080504.
36. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315:629-34. doi: 10.1136/bmj.315.7109.629.
37. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics*. 1994;50:1088-101. doi: 10.2307/2533446.
38. Wang A, Zhu C, Fu L, Wan X, Yang X, Zhang H, Miao R, He L, Sang X, Zhao H. Citrus Fruit intake substantially reduces the risk of esophageal cancer: a meta-analysis of epidemiologic studies. *Medicine (Baltimore)*. 2015;94:e1390. doi: 10.1097/md.0000000000001390.
39. Bravi F, Bosetti C, Filomeno M, Levi F, Garavello W, Galimberti S, Negri E, La Vecchia C. Foods, nutrients and the risk of oral and pharyngeal cancer. *Br J Cancer*. 2013;109:2904-10. doi: 10.1038/bjc.2013.667.
40. Bae JM, Lee EJ, Guyatt G. Citrus fruit intake and pancreatic cancer risk: a quantitative systematic review. *Pancreas*. 2009;38:168-74. doi: 10.1097/MPA.0b013e318188c497.
41. Song JK, Bae JM. Citrus fruit intake and breast cancer risk: a quantitative systematic review. *J Breast Cancer*. 2013;16:72-6. doi: 10.4048/jbc.2013.16.1.72.
42. Bo Y, Lu Y, Zhao Y, Zhao E, Yuan L, Lu W, Cui L, Lu Q. Association between dietary vitamin C intake and risk of esophageal cancer: A dose-response meta-analysis. *Int J Cancer*. 2016;138:1843-50. doi: 10.1002/ijc.29838.
43. Lee KW, Lee HJ, Surh YJ, Lee CY. Vitamin C and cancer chemoprevention: reappraisal. *Am J Clin Nutr*. 2003;78:1074-8. doi: 10.1093/ajcn/78.6.1074.
44. Palozza P. Carotenoids and modulation of cancer: Molecular targets. *Current Pharmacogenomics*. 2004;2:35-45. doi: 10.2174/1570160043476169.
45. Zeng FF, Liu YT, Lin XL, Fan YY, Zhang XL, Xu CH, Chen YM. Folate, vitamin B6, vitamin B12 and methionine intakes and risk for nasopharyngeal carcinoma in Chinese adults: a matched case-control study. *Br J Nutr*. 2016;115:121-8. doi: 10.1017/s0007114515004146.
46. Bayazit V, Konar V. Biochemical and physiological evaluations of limonoids as potential cancer destroyers. *Journal of Animal and Veterinary Advances*. 2010;9:1099-107. doi: 10.3923/javaa.2010.1099.1107.
47. Perez JL, Jayaprakasha GK, Cadena A, Martinez E, Ahmad H, Patil BS. In vivo induction of phase II detoxifying enzymes, glutathione transferase and quinone reductase by citrus triterpenoids. *BMC Complement Altern Med*. 2010;10: 51. doi: 10.1186/1472-6882-10-51.
48. Hung WL, Suh JH, Wang Y. Chemistry and health effects of furanocoumarins in grapefruit. *J Food Drug Anal*. 2017;25:71-83. doi: 10.1016/j.jfda.2016.11.008.
49. Feng BJ, Huang W, Shugart YY, Lee MK, Zhang F, Xia JC et al. Genome-wide scan for familial nasopharyngeal carcinoma reveals evidence of linkage to chromosome 4. *Nat Genet*. 2002;31:395-9. doi: 10.1038/ng932.
50. Ding M, Ellervik C, Huang T, Jensen MK, Curhan GC, Pasquale LR et al. Diet quality and genetic association with body mass index: results from 3 observational studies. *Am J Clin Nutr*. 2018;108:1291-300. doi: 10.1093/ajcn/nqy203.
51. Rottenberg Y, Levine H, Keinan-Boker L, Derazne E, Leiba A, Kark JD. Risk of nasopharyngeal carcinoma penetrates across immigrant generations: A migrant cohort study of 2.3 million Jewish Israeli adolescents. *Int J Cancer*. 2017;140:1060-7. doi: 10.1002/ijc.30525.
52. Roy Chattopadhyay N, Das P, Chatterjee K, Choudhuri T. Higher incidence of nasopharyngeal carcinoma in some regions in the world confers for interplay between genetic factors and external stimuli. *Drug Discov Ther*. 2017;11:170-80. doi: 10.5582/ddt.2017.01030.
53. Iwase Y, Takemura Y, Ju-ichi M, Kawaii S, Yano M, Okuda Y et al. Inhibitory effect of Epstein-Barr virus activation by Citrus fruits, a cancer chemopreventor. *Cancer Lett*. 1999;139:227-36. doi: 10.1016/s0304-3835(99)00041-5.
54. Tsao SW, Yip YL, Tsang CM, Pang PS, Lau VM, Zhang G, Lo KW. Etiological factors of nasopharyngeal carcinoma. *Oral Oncol*. 2014;50:330-8. doi: 10.1016/j.oraloncology.2014.02.006.
55. Swan SH, Shaw GM, Schulman J. Reporting and selection bias in case-control studies of congenital malformations. *Epidemiology*. 1992;3:356-63. doi: 10.1097/00001648-199207000-00011.
56. Stephanie M G. Fruit and vegetable intake and risk of cancer: a prospective cohort study. *Am J Clin Nutr*. 2009;89:347-53. doi: 10.3945/ajcn.2008.26722.



Supplemental figure 1. Funnel plot of the relative risk (for highest-versus-lowest intake category) versus the standard error of the log relative risk for studies evaluating citrus fruit intake and nasopharyngeal carcinoma (NPC).

Supplementary table 1. Characteristics of studies no having OR or RR for citrus fruit in Figure 1

First author, year	Country	Study design	Case/Subjects (n)	Sex (%male)	Age (years)	Exposure assessment	Diagnosis method	Adjustments
Yu et al, 1986	China	Case-control (PCC)	250/500	64.0	<35	Interview	Medical records	Age, sex
Ning et al, 1990	China	Case-control (PCC)	100/400	68.0	≤64	Interview	Pathology reports	Age, sex, race
Lee et al, 1994	Singapore	Case-control (HCC)	200/606	71.7	NR	Interview	Medical records	Age, sex, education, dialect group
Cheng et al, 1999	China	Case-control (PCC)	375/702	68.8	<75	Interview	Pathology reports	Age, sex, race, educational level, family history of NPC
Shahar et al, 2004	Malaysia	Case-control (HCC)	100/200	50.0	≥18	Interview	Medical records	Age, sex, ethnic origin
Ekburanawat et al, 2010	Thailand	Case-control (PCC)	327/654	74.0	48.1 (mean)	Interview	Pathology reports	Sex, age, geographic residence, cigarettes smoking, education
Jia et al, 2010	China	Case-control (HCC)	1387/2846	72.5	<80	Interview	Pathology reports	Age, sex, education, dialect, household type
Turkoz et al, 2011	Turkey	Case-control (PCC&HCC)	183/366	65.6	18-76	Interview	Pathology reports	Sex, age
Fachiroh et al, 2012	Thailand	Case-control (PCC)	681/1759	66.1	48.0 (mean)	Interview	Pathology reports	Sex, age group, center, education, alcohol drinking, smoking status
Hashim et al, 2012	Malaysia	Case-control (HCC)	48/96	81.3	47.5 (mean)	Interview	Medical records	Sex, age, ethnicity
Shen et al, 2012	Italy	Cohort	1533	73.3	46.1 (mean)	Interview	Medical records	Age, BMI, spouse, education, clinical stage, smoking status, alcohol intake
Belbaraka et al, 2013	Morocco	Case-control	60/120	60.0	≥25	Interview	Medical records	Age, sex, socio economic level
Yong et al, 2017	Singapore	Case-control (PCC)	290/580	81.4	21-80	Interview	Pathology reports	Sex, ethnicity, age, educational level

BMI: Body Mass Index; HCC: hospital-based case-control study; NPC: nasopharyngeal carcinoma; NR: not reported; OR: odds ratio; PCC: population-based case-control study; RR: relative risk.

Supplementary table 2. Citrus fruit intake in association with NPC risk in each original study

First author, year	Comparison of exposure level	OR (95% CI)
Chen et al, 1997 ¹⁸	<Monthly	1.00
	<weekly	NR
	1-4 times/week	NR
	>4 times/week	0.66 (0.50-0.89)
Farrow et al, 1998 ¹⁹	Quartile 1	1.00
	Quartile 2	1.02 (0.53-1.98)
	Quartile 3	0.46 (0.22-0.96)
	Quartile 4	0.68 (0.33-1.41)
Armstrong et al, 1998 ²⁰	≤Monthly	1.00
	<weekly	0.40 (0.24-0.69)
	≥weekly	0.52 (0.31-0.85)
Yuan et al, 2000 ²¹	<Monthly	1.00
	Monthly	0.71 (0.53-0.96)
	Weekly	0.55 (0.41-0.73)
	Daily	0.55 (0.38-0.78)
Feng et al, 2007 ²²	<Monthly	1.00
	<3 times/week	1.40 (0.80-2.20)
	≥3 times/week	1.50 (0.90-2.60)
Nešić et al, 2010 ²³	<Monthly	1.00
	≤3 times /week	0.29 (0.06-1.36)
Liu et al, 2012 ²⁴	Quartile 1	1.00
	Quartile 2	0.95 (0.67-1.35)
	Quartile 3	0.74 (0.50-1.08)
	Quartile 4	0.65 (0.43-0.96)
Hsu et al, 2012 ²⁵	≤2 times/week	1.00
	2-6 times/week	1.06 (0.69-1.61)
	>6 times/week	1.03 (0.71-1.51)
Polesel et al, 2013 ²⁶	<weekly	1.00
	1-4 times/week	0.59 (0.33-1.05)
	4-7 times/week	0.96 (0.60-1.52)
	>7 times/week	0.75 (0.45-1.26)

CI: confidence interval; NPC: nasopharyngeal carcinoma; NR: not reported; OR: odds ratio.

Supplementary Table 3. Sensitivity analysis of the association between citrus fruit intake and NPC

First author, year	Pooled OR	95% CI
Chen et al, 1997 ¹⁸	0.74	0.56, 0.97
Farrow et al, 1998 ¹⁹	0.73	0.57, 0.93
Armstrong et al, 1998 ²⁰	0.75	0.59, 0.96
Yuan et al, 2000 ²¹	0.76	0.59, 0.97
Feng et al, 2007 ²²	0.67	0.57, 0.80
Nešić et al, 2010 ²³	0.74	0.59, 0.92
Liu et al, 2012 ²⁴	0.74	0.57, 0.96
Hsu et al, 2012 ²⁵	0.68	0.54, 0.86
Polesel et al, 2013 ²⁶	0.72	0.56, 0.93

CI: confidence interval; NPC: nasopharyngeal carcinoma; OR: odds ratio.