

This author's PDF version corresponds to the article as it appeared upon acceptance. Fully formatted PDF versions will be made available soon.

Weight management for adolescents with overweight and obesity: current treatment

doi: 10.6133/apjcn.202509/PP.0002

Published online: September 2025

Running title: Weight management in adolescents

Liru Chen MD¹, Xingyu Zhu², Chengyu Liu MM³

¹Department of Nutrition, Beijing Hospital, National Center of Gerontology, Institute of Geriatric Medicine, Chinese Academy of Medical Sciences, Beijing, China

²The high school Affiliated to Renmin University of China, Beijing, China

³Department of General Surgery, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China

Authors' email addresses and contributions:

LC: chenliruabc@163.com

Contribution: conceived the study question, and contributed to the study design, and writing the manuscript.

XZ: zxy13681323537@163.com

Contribution: conceived the study question, collecting and organizing literature, and writing the manuscript.

CL: chengyujoe@gmail.com

Contribution: collecting and organizing literature, and revising the manuscript critically for important content..

Corresponding Author: Dr Liru Chen, Department of Nutrition, Beijing Hospital, National Center of Gerontology, Institute of Geriatric Medicine, Chinese Academy of Medical Sciences, Beijing, China. Tel: +86 15901116313. Email: chenliruabc@163.com

ABSTRACT

Background and Objectives: The prevalence of overweight or obesity in adolescents is steadily increasing in most countries around the world. Adolescent obesity increases the risk of chronic diseases such as cardiovascular disease, and is associated with negative health consequences, increasing the burden on health services. **Methods and Study Design:** Literature searching was conducted in PubMed and Google Scholar using the keywords “overweight”, “obesity”, “adolescent”, “weight management”, “dietary management” and “nutritional intervention” combined with Boolean operators “AND” and “OR”. **Results:** Multicomponent lifestyle interventions, including diet, physical activity, and behavioral interventions, are used as first-line treatment for anti-obesity interventions. Dietary management methods such as energy-restricted diet are beneficial to control body weight, and it is important to ensure their normal growth and development while restricting energy. **Conclusions:** Multicomponent lifestyle intervention is the first choice for anti-obesity intervention. It is recommended that intensive health behaviour lifestyle treatment combined with anti-obesity medications be used at the beginning of anti-obesity treatment in adolescents.

Key Words: weight management, overweight, obesity, adolescent, multicomponent lifestyle intervention

INTRODUCTION

A recent meta-analysis revealed that the worldwide prevalence of obesity in children and adolescents was 8.5% (95% CI, 8.2 to 8.8) and the prevalence of overweight in adolescents was 22.2% (95% CI, 21.6 to 22.8).¹ From 1990 to 2022, the global prevalence of obesity among adolescents has increased.^{2, 3} The estimated number of girls and boys with obesity in 2022 was 65.1 million (95% CI, 59.4 to 71.7) and 94.2 million (95% CI, 85.3 to 103.0), respectively, an increase of 51.2 million (95% CI, 45.2 to 57.8) and 76.7 million (95% CI, 67.6 to 85.7), respectively, since 1990.² Obesity in adolescents is defined as body mass index (BMI) more than 2 standard deviations above the median world health organization (WHO) growth reference value.⁴ The prevalence of obesity among 12- to 19-year-old in the United States has increased steadily, from 16.8% (95% CI, 14.2 to 19.8) in 2007-2008 to 18.5% (95% CI, 15.8 to 21.3) in 2015-2016.^{3, 5, 6} In China, the 2015 national survey revealed that the prevalence of obesity in children and adolescents aged 7-17 years was 13.2% (95%CI, 12.8 to 13.7).⁷

It should be emphasized that adolescent obesity is a chronic condition requiring lifelong care.^{8, 9} Clinical evidence shows that persistent obesity in adolescents increases the risk of chronic diseases such as cardiovascular disease, hypertension, diabetes mellitus and future cancer.¹⁰⁻¹⁴ Based on the data from the global burden of disease collaborative for 2000 and 2013, it is estimated that by 2025 global childhood and adolescent obesity will lead to 12 million children and adolescents with impaired glucose tolerance, 4 million with type 2 diabetes, 27 million with hypertension and 38 million with hepatic steatosis.¹⁵ There is also a growing recognition that adolescent obesity is associated with a range of immediate and long-term negative health consequences, reduced quality of life, increased economic costs for individuals and society, and increased burden on health services.^{16, 17} In addition, the adolescent period is characterized by the development of autonomy and self-identity, and the adolescents with overweight and obesity is prone to low self-esteem, depression and anxiety, which will affect mental health status and quality of life.^{1, 16, 18}

Lifestyle interventions, including dietary management, play an important role in the treatment of adolescents overweight and obesity, and recent clinical studies have provided reliable evidence of the meaningful reduction in body weight of anti-obesity medications.^{8, 19} This review focuses on the risk factors and pathophysiology of overweight and obesity in adolescents, as well as approaches to its treatment for weight management.

MATERIALS AND METHODS

Literature searching was conducted in PubMed and Google Scholar using the keywords “overweight”, “obesity”, “adolescent”, “weight management”, “dietary management” and “nutritional intervention” combined with Boolean operators “AND” and “OR”.

This review included peer-reviewed clinical trials, systematic reviews, and meta-analyses published in English that evaluated the effects of anti-obesity treatments such as dietary interventions, exercise interventions, anti-obesity medications, and metabolic surgery on weight management in adolescents with overweight or obesity.

RESULTS

Risk factors and possible pathogenesis of overweight and obesity in adolescents

Unlike childhood obesity, which is linked to many environments in early life, genetic, lifestyle, behavioural, social, and environmental factors are major risk factors for obesity in adolescents.^{20, 21} Obesity, dominated by polygenic variants, is the result of interactions between genetic predisposition and environmental factors.²² Many lifestyle and behavioural

factors increase the risk of obesity, including reduced physical activity, being sedentary, short and poor sleep duration, fast food, carbonated soft drink intake, and no daily vegetable intake.²³⁻²⁶ Social and environmental factors include chronic psychosocial stress, widespread marketing of food and beverages to adolescents, a shift from conventional to ultra-processed foods, loss of walkable green spaces in many urban settings, increased motorized transportation, rapid changes in the use of technology, and environmental contaminants, et al.^{20, 27, 28} Individual, social, and environmental risk factors often overlap and/or influence each other and may have longitudinal effects throughout adolescence, leading to weight gain and exacerbating existing obesity.⁸

The simple explanatory model of obesity (too much caloric intake and too little physical activity) has been replaced by a complex multi-component model that includes genetics, social and cultural backgrounds, obesogenic environment, and the barriers to treatment.⁹ Body weight is primarily regulated by two systems: energy homeostasis and cognitive-emotional control, and obesity occurs when genetic and epigenetic factors, behavioural risk patterns, and broader social and environmental factors influence both systems.^{20, 29} One is a homeostatic system operating at an unconscious level regulated primarily by brain centres in the hypothalamus and brainstem, receiving long-term signals from energy stores in adipose tissue (such as leptin) and short-term hunger and satiety signals from the gastrointestinal tract.^{20, 30} Another non-homeostatic system (cognition-emotion control) is regulated by higher brain centres and operates at the level of consciousness, when homeostatic signals are integrated in the thalamus, combined with stimuli from the environment, experience and emotion, and then induced emotional and cognitive impulses to control food intake.^{20, 29, 31}

The above risk factors and possible pathogenesis are summarized in Table 1. Meanwhile, Table 2 summarizes the interventions and strategies for adolescents with overweight and obesity.

Dietary management in adolescents with overweight and obesity

Obesity is a chronic disease that should be treated using intensive and long-term care strategies that provide continuous anti-obesity treatment and medical monitoring, a key component of anti-obesity treatment is simultaneous monitoring and treatment of comorbidities, which are not discussed in our review.⁸ Nutritional therapy including diet management is the preferred treatment for overweight and obesity in adolescents, and it is important to ensure their normal growth and development while restricting energy.³² Medical nutrition therapy aims to achieve positive health outcomes in addition to weight management,

and obesity-related complications and other comorbidities should be managed independently of any anti-obesity treatment.³³ A recent meta-analysis suggests that obesity interventions that include dietary components significantly reduce average total energy intake, increase fruit and/or vegetable intake, and reduce the consumption of sugar-sweetened beverages in adolescents with overweight or obesity.³⁴

Energy restriction diet is beneficial for overweight and obese adolescents to control body weight, and reduce energy supply on the premise of ensuring the energy required for normal growth and development, and prevent any impact upon growth under the supervision of a health professional such as a dietitian.^{32, 35} To reduce body weight, dietary interventions should be provided by nutrition professionals to produce an energy deficit of 500 to 750 kcal per day.³³ Structured meal plans, portion control, and meal replacements as possible weight management interventions for adolescents, and the Mediterranean diet, vegetarian diet, the Dietary Approaches to Stop Hypertension, and low-carbohydrate diet have been shown to be associated with improved metabolic health, with or without weight change.^{33, 35} The current guidelines do not emphasize the dietary composition of nutritional therapy, but focus on balanced and healthy diets.³³ The unique characteristics of adolescent development can influence food choices, and adolescents should be actively involved in developing healthy dietary patterns.³⁶ Recent studies have focused on intermittent energy restriction as a treatment option for obesity, but there is no evidence that intermittent energy restriction has significant advantages over continuous energy restriction.^{33, 37, 38} Very low-calorie diets (calorie intake \leq 800 kcal/ day) are not recommended for the treatment of obesity but only for certain medical conditions, which always requires medical supervision.^{32, 33}

Other types of dietary management have also been explored in several clinical studies. The 24-week low-energy anti-inflammatory diet intervention significantly reduced body weight and visceral adipose tissue, and improved cardiometabolic and inflammatory status in obese patients.³⁹ A randomized controlled trial did not observe a significant effect of a high protein/low glycemic index diet intervention versus a moderate protein/moderate glycemic index diet intervention on insulin resistance in overweight/obese adolescents.⁴⁰ A meta-analysis showed that dietary management were associated with a reduction in the prevalence of eating disorders, risk of eating disorders, and symptoms in children and adolescents with overweight or obesity.⁴¹ When using diet management for overweight and obesity, pediatricians should also assess and address modifiable risk factors for obesity, including the frequency of eating out and fast-food consumption.^{42, 43}

Physical activity and behavioural management in overweight or obese adolescents

In addition to dietary intervention, comprehensive lifestyle interventions include physical activity and behavioural intervention.^{8, 33, 44} Increasing daily activities and physical exercise are beneficial to increase energy expenditure in adolescents with overweight and obesity, which combined with dietary intervention can keep energy metabolism in a negative balance.⁴⁵ It is recommended 30 min moderate-intensity aerobic exercise at least 5 days or more per week, combined with strength or resistance training.⁴⁵ In children and adolescents with moderate-to-severe obesity, a 16-week exercise intervention significantly reduced percentage body fat and cardiometabolic risk markers and increased lean mass and leg muscle strength compared with a usual-care multidisciplinary lifestyle intervention program.⁴⁶ There is high-quality study revealed that physical activity interventions improve average cognitive executive function scores in children and adolescents with overweight or obesity.⁴⁷ Play-based physical activity and nutrition advice interventions can improve quality of life in overweight and obese children.⁴⁸ Short-term low-volume high-intensity interval training is an effective way to improve the cardiorespiratory fitness of adolescents, which suggests that participation in physical education classes in the school curriculum has health benefits.⁴⁹ Exercise interventions targeting special groups such as those with comorbid asthma or psychological disorders require additional adjustments, increased control of underlying diseases, family support, and social support.^{50, 51}

Multicomponent lifestyle interventions, including diet, physical activity, and behaviour modification, are used as first-line treatment for anti-obesity interventions.³⁵ Intensive healthy behaviour and lifestyle treatment is an essential approach to promote the maintenance of healthier eating and physical activity habits to achieve weight management in adolescents, in which patients and families cooperate with multidisciplinary treatment teams to conduct health education and skill training on multiple topics, behaviour change and counselling.^{8, 52, 53} The main views of adolescents who had participated in anti-obesity interventions were "fear of receiving life intervention before intervention", "hope for long-term support", "enjoy sports and physical activity", and "strong motivation to improve body image and social goodwill", and consideration of the above perspectives may help policy makers to formulate future interventions.⁵⁴ Intensive behavioural intervention programs can reduce BMI, reduce comorbidities, and improve quality of life in children and adolescents.^{8, 55, 56} Comprehensive, intensive behavioural interventions (≥ 26 h of contact) in obese children and adolescents aged 6 years and older can improve weight status for up to 12 months.⁵⁷

A multi-centre study in India showed that school-based lifestyle intervention was more effective in improving physical activity, lifestyle behaviours and quality of life in adolescents than exercise intervention alone.^{58, 59} A motivation-based educational program is more effective than dietary counselling in reducing wrist circumference and improving quality of life among adolescents with overweight and obesity.⁶⁰ Parents and caregivers can play a key role in the treatment of obesity through strategies such as parental supervision and changing the family environment, and studies have found that family-based interventions have a significant indirect effect on the treatment of obesity through the quality of parent-child relationship and dietary attitudes.^{61, 62} Current evidence suggests that behaviour-oriented prevention programs have limited long-term effects, and behaviour-based programs can be combined with community-based prevention to combat "obesogenic environments".^{20, 63}

Anti-obesity medications and metabolic surgery

The new clinical practice guideline recommends that anti-obesity medications therapy be provided to adolescents with obesity in parallel with intensive health behaviour lifestyle treatment, which represents a paradigm shift from pharmacotherapy when lifestyle therapy fails to a combination of intensive health behaviour lifestyle treatment and anti-obesity medications from the start of treatment.^{8, 9}

A recent meta-analysis provides evidence that anti-obesity medications reduce BMI (-1.71, 95% CI: -2.27 to -1.14), but may require dose modifications, with 1 in 100 adolescents experiencing serious adverse events, with semaglutide achieving the greatest reduction in BMI (-5.88, 95% CI: -6.99 to -4.77).¹⁹ Another meta-analysis provided solid evidence for the effect of glucagon-like peptide-1 receptor agonists (GLP1Ras) on weight loss in non-diabetic individuals with overweight or obesity, with significant weight loss (MD -5.319 kg, 95% CI: -6.465 -- 4.174), BMI (MD -2.373 kg/m², 95% CI: -2.821 to 1.924), and waist circumference (MD - 4.302 cm, 95% CI :-5.185 to 3.419).⁶⁴

In overweight or obese participants, semaglutide plus lifestyle intervention was associated with weight loss and a greater reduction in abdominal visceral fat area, suggesting that this is a promising treatment option for weight management.^{65, 66} A phase III clinical trial found that liraglutide significantly reduced BMI in obese adolescents.^{67, 68} A randomized controlled trial showed that orlistat improved liver enzymes, body weight, BMI, waist circumference, total cholesterol, and low-density lipoprotein cholesterol in overweight/obese adolescents with non-alcoholic fatty liver disease.⁶⁹ An economic evaluation study of quality-adjusted life-years, costs, and incremental cost-effectiveness ratios for anti-obesity medications suggested

that the top-dose phentermine and topiramate assisted lifestyle counselling was estimated to be cost-effective after 5 years.⁷⁰

Anti-obesity medications may play a role in combating the pathophysiology of the development of obesity in adolescents through the following mechanisms. GLP1Ras reduce appetite by acting on hypothalamic glucagon-like peptide-1 receptors, increasing satiety and reducing food reward and craving by activating the hindbrain and slowing gastric emptying.⁷¹ Phentermine reduces appetite by inhibiting reuptake of norepinephrine, and topiramate reduces caloric intake by increasing gamma-aminobutyric acid activity and inhibiting carbonic anhydrase.⁷¹

Relevant clinical guidelines recommend that metabolic and bariatric surgery be considered for youth with class II obesity (defined as a BMI of 120% of the 95th percentile) or class III obesity (defined as a BMI of 140% of the 95th percentile) with comorbidities such as type 2 diabetes and obstructive sleep apnea.^{8, 72} Metabolic and bariatric surgery includes adjustable gastric band, sleeve gastrectomy, Roux-en-Y gastric bypass and biliopancreatic diversion/duodenal switch.⁷³ After bariatric surgery, long-term multidisciplinary follow-up care is required, including diet and micronutrient monitoring, individualized nutritional supplementation and psychological support.³³

Conclusion

The prevalence of adolescent obesity has been steadily increasing in most countries around the world, and adolescent obesity increases the risk of chronic diseases such as cardiovascular disease. Multicomponent lifestyle interventions, including diet, physical activity, and behavioural interventions, are used as first-line treatment for anti-obesity interventions. It is important to maintain normal growth and development on an energy-restricted diet, while increasing daily activity and physical exercise is also necessary. Clinical practice guidelines recommend combining intensive health behaviour lifestyle treatment with anti-obesity medications at the beginning of anti-obesity treatment in adolescents. A variety of anti-obesity medications can significantly reduce body weight and BMI, but serious adverse events should be alert. Metabolic bariatric surgery should strictly follow the indications of clinical guidelines, and long-term multidisciplinary follow-up care is needed.

CONFLICT OF INTEREST AND FUNDING DISCLOSURE

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

This research was supported by China International Medical Foundation [Z-2017-24-2211] and Jinqiao Project of Beijing Association for Science and Technology [ZZ22058].

REFERENCES

1. Zhang X, Liu J, Ni Y, Yi C, Fang Y, Ning Q, et al. Global Prevalence of Overweight and Obesity in Children and Adolescents: A Systematic Review and Meta-Analysis. *JAMA Pediatr.* 2024;178(8):800-13.
2. (NCD-RisC) NRFC. Worldwide trends in underweight and obesity from 1990 to 2022: a pooled analysis of 3663 population-representative studies with 222 million children, adolescents, and adults. *Lancet.* 2024;403(10431):1027-50.
3. Hales CM, Fryar CD, Carroll MD, Freedman DS, Ogden CL. Trends in Obesity and Severe Obesity Prevalence in US Youth and Adults by Sex and Age, 2007-2008 to 2015-2016. *JAMA.* 2018;319(16):1723-5.
4. de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ.* 2007;85(9):660-7.
5. Stierman B, Afful J, Carroll MD, Chen TC, Davy O, Fink S, et al. National Health and Nutrition Examination Survey 2017-March 2020 Prepandemic Data Files-Development of Files and Prevalence Estimates for Selected Health Outcomes. *Natl Health Stat Report.* 2021(158).
6. Hales CM, Carroll MD, Fryar CD, Ogden CL. Prevalence of Obesity Among Adults and Youth: United States, 2015-2016. *NCHS Data Brief.* 2017(288):1-8.
7. Zhou Y, Zhang Q, Wang T, Zhang Y, Xu B. Prevalence of overweight and obesity in Chinese children and adolescents from 2015. *Ann Hum Biol.* 2017;44(7):642-3.
8. Hampl SE, Hassink SG, Skinner AC, Armstrong SC, Barlow SE, Bolling CF, et al. Clinical Practice Guideline for the Evaluation and Treatment of Children and Adolescents With Obesity. *Pediatrics.* 2023;151(2).
9. Reinehr T, Dieris B. New clinical practice guideline for evaluation and treatment of children and adolescents with obesity: paradigm shifts. *Lancet Diabetes Endocrinol.* 2023;11(4):222-3.
10. Twig G, Yaniv G, Levine H, Leiba A, Goldberger N, Derazne E, et al. Body-Mass Index in 2.3 Million Adolescents and Cardiovascular Death in Adulthood. *N Engl J Med.* 2016;374(25):2430-40.
11. Twig G, Zucker I, Afek A, Cukierman-Yaffe T, Bendor CD, Derazne E, et al. Adolescent Obesity and Early-Onset Type 2 Diabetes. *Diabetes Care.* 2020;43(7):1487-95.
12. Xu T, Zhu G, Liu J, Han S. Gender-specific prevalence and associated risk factors of high normal blood pressure and hypertension among multi-ethnic Chinese adolescents aged 8-18 years old. *Blood Press.* 2015;24(3):189-95.
13. Huh Y, Park HS. Associations of overweight and obesity with cardiometabolic risk factor clusters among Korean adolescents. *Sci Rep.* 2024;14(1):3581.

14. Furer A, Afek A, Sommer A, Keinan-Boker L, Derazne E, Levi Z, et al. Adolescent obesity and midlife cancer risk: a population-based cohort study of 2.3 million adolescents in Israel. *Lancet Diabetes Endocrinol.* 2020;8(3):216-25.
15. Lobstein T, Jackson-Leach R. Planning for the worst: estimates of obesity and comorbidities in school-age children in 2025. *Pediatr Obes.* 2016;11(5):321-5.
16. Griffiths LJ, Parsons TJ, Hill AJ. Self-esteem and quality of life in obese children and adolescents: a systematic review. *Int J Pediatr Obes.* 2010;5(4):282-304.
17. Ling J, Chen S, Zahry NR, Kao TA. Economic burden of childhood overweight and obesity: A systematic review and meta-analysis. *Obes Rev.* 2023;24(2):e13535.
18. Hadjiyannakis S, Ibrahim Q, Li J, Ball GDC, Buchholz A, Hamilton JK, et al. Obesity class versus the Edmonton Obesity Staging System for Pediatrics to define health risk in childhood obesity: results from the CANPWR cross-sectional study. *Lancet Child Adolesc Health.* 2019;3(6):398-407.
19. Torbahn G, Jones A, Griffiths A, Matu J, Metzendorf MI, Ells LJ, et al. Pharmacological interventions for the management of children and adolescents living with obesity-An update of a Cochrane systematic review with meta-analyses. *Pediatr Obes.* 2024;19(5):e13113.
20. Lister NB, Baur LA, Felix JF, Hill AJ, Marcus C, Reinehr T, et al. Child and adolescent obesity. *Nat Rev Dis Primers.* 2023;9(1):24.
21. Bouchard C. Genetics of Obesity: What We Have Learned Over Decades of Research. *Obesity* (Silver Spring). 2021;29(5):802-20.
22. Hinney A, Körner A, Fischer-Posovszky P. The promise of new anti-obesity therapies arising from knowledge of genetic obesity traits. *Nat Rev Endocrinol.* 2022;18(10):623-37.
23. Farooq A, Martin A, Janssen X, Wilson MG, Gibson AM, Hughes A, et al. Longitudinal changes in moderate-to-vigorous-intensity physical activity in children and adolescents: A systematic review and meta-analysis. *Obes Rev.* 2020;21(1):e12953.
24. Mahumud RA, Sahle BW, Owusu-Addo E, Chen W, Morton RL, Renzaho AMN. Association of dietary intake, physical activity, and sedentary behaviours with overweight and obesity among 282,213 adolescents in 89 low and middle income to high-income countries. *Int J Obes (Lond).* 2021;45(11):2404-18.
25. Fatima Y, Doi SA, Mamun AA. Sleep quality and obesity in young subjects: a meta-analysis. *Obes Rev.* 2016;17(11):1154-66.
26. Calcaterra V, Rossi V, Tagi VM, Baldassarre P, Grazi R, Taranto S, et al. Food Intake and Sleep Disorders in Children and Adolescents with Obesity. *Nutrients.* 2023;15(22).
27. Ford ND, Patel SA, Narayan KM. Obesity in Low- and Middle-Income Countries: Burden, Drivers, and Emerging Challenges. *Annu Rev Public Health.* 2017;38:145-64.
28. Neufeld LM, Andrade EB, Ballonoff Suleiman A, Barker M, Beal T, Blum LS, et al. Food choice in transition: adolescent autonomy, agency, and the food environment. *Lancet.* 2022;399(10320):185-97.
29. Berthoud HR, Münzberg H, Morrison CD. Blaming the Brain for Obesity: Integration of Hedonic and Homeostatic Mechanisms. *Gastroenterology.* 2017;152(7):1728-38.

30. Berthoud HR, Morrison CD, Münzberg H. The obesity epidemic in the face of homeostatic body weight regulation: What went wrong and how can it be fixed? *Physiol Behav.* 2020;222:112959.
31. Devoto F, Zapparoli L, Bonandrini R, Berlingeri M, Ferrulli A, Luzi L, et al. Hungry brains: A meta-analytical review of brain activation imaging studies on food perception and appetite in obese individuals. *Neurosci Biobehav Rev.* 2018;94:271-85.
32. Guidelines for medical nutrition treatment of overweight/obesity in China (2021). *Asia Pac J Clin Nutr.* 2022;31(3):450-82.
33. Hassapidou M, Vlassopoulos A, Kalliostra M, Govers E, Mulrooney H, Ells L, et al. European Association for the Study of Obesity Position Statement on Medical Nutrition Therapy for the Management of Overweight and Obesity in Adults Developed in Collaboration with the European Federation of the Associations of Dietitians. *Obes Facts.* 2023;16(1):11-28.
34. Duncanson K, Shrewsbury V, Burrows T, Chai LK, Ashton L, Taylor R, et al. Impact of weight management nutrition interventions on dietary outcomes in children and adolescents with overweight or obesity: a systematic review with meta-analysis. *J Hum Nutr Diet.* 2021;34(1):147-77.
35. Alman KL, Lister NB, Garnett SP, Gow ML, Aldwell K, Jebeile H. Dietetic management of obesity and severe obesity in children and adolescents: A scoping review of guidelines. *Obes Rev.* 2021;22(1):e13132.
36. Roberts KJ, Hafez SA, Snethen J, Binns HJ, Knafl KA. Perceptions of Weight Management: Interviews with Adolescents with Severe Obesity and Their Mothers. *Child Obes.* 2022;18(4):219-27.
37. Carter S, Clifton PM, Keogh JB. Effect of Intermittent Compared With Continuous Energy Restricted Diet on Glycemic Control in Patients With Type 2 Diabetes: A Randomized Noninferiority Trial. *JAMA Netw Open.* 2018;1(3):e180756.
38. Li M, Li J, Xu Y, Gao J, Cao Q, Ding Y, et al. Effect of 5:2 Regimens: Energy-Restricted Diet or Low-Volume High-Intensity Interval Training Combined With Resistance Exercise on Glycemic Control and Cardiometabolic Health in Adults With Overweight/Obesity and Type 2 Diabetes: A Three-Arm Randomized Controlled Trial. *Diabetes Care.* 2024;47(6):1074-83.
39. Kendel Jovanović G, Mrakovcic-Sutic I, Pavičić Žeželj S, Šuša B, Rahelić D, Klobučar Majanović S. The Efficacy of an Energy-Restricted Anti-Inflammatory Diet for the Management of Obesity in Younger Adults. *Nutrients.* 2020;12(11).
40. Dorenbos E, Drummen M, Adam T, Rijks J, Winkens B, Martínez JA, et al. Effect of a high protein/low glycaemic index diet on insulin resistance in adolescents with overweight/obesity-A PREVIEW randomized clinical trial. *Pediatr Obes.* 2021;16(1):e12702.
41. Jebeile H, Gow ML, Baur LA, Garnett SP, Paxton SJ, Lister NB. Treatment of obesity, with a dietary component, and eating disorder risk in children and adolescents: A systematic review with meta-analysis. *Obes Rev.* 2019;20(9):1287-98.
42. Nago ES, Lachat CK, Dossa RA, Kolsteren PW. Association of out-of-home eating with anthropometric changes: a systematic review of prospective studies. *Crit Rev Food Sci Nutr.* 2014;54(9):1103-16.

43. Jiang Y, Wang J, Wu S, Li N, Wang Y, Liu J, et al. Association between Take-Out Food Consumption and Obesity among Chinese University Students: A Cross-Sectional Study. *Int J Environ Res Public Health*. 2019;16(6).
44. Kang E, Hong YH, Kim J, Chung S, Kim KK, Haam JH, et al. Obesity in Children and Adolescents: 2022 Update of Clinical Practice Guidelines for Obesity by the Korean Society for the Study of Obesity. *J Obes Metab Syndr*. 2024;33(1):11-9.
45. Semlitsch T, Stigler FL, Jeitler K, Horvath K, Siebenhofer A. Management of overweight and obesity in primary care-A systematic overview of international evidence-based guidelines. *Obes Rev*. 2019;20(9):1218-30.
46. Seo YG, Lim H, Kim Y, Ju YS, Lee HJ, Jang HB, et al. The Effect of a Multidisciplinary Lifestyle Intervention on Obesity Status, Body Composition, Physical Fitness, and Cardiometabolic Risk Markers in Children and Adolescents with Obesity. *Nutrients*. 2019;11(1).
47. Martin A, Booth JN, Laird Y, Sproule J, Reilly JJ, Saunders DH. Physical activity, diet and other behavioural interventions for improving cognition and school achievement in children and adolescents with obesity or overweight. *Cochrane Database Syst Rev*. 2018;3(3):Cd009728.
48. Aguilar-Cordero MJ, León Ríos XA, Rojas-Carvajal AM, Latorre-García J, Expósito-Ruiz M, Sánchez-López AM. Effects of physical activity on quality of life in overweight and obese children. *Nutr Hosp*. 2021;38(4):736-41.
49. Martin-Smith R, Cox A, Buchan DS, Baker JS, Grace F, Sculthorpe N. High Intensity Interval Training (HIIT) Improves Cardiorespiratory Fitness (CRF) in Healthy, Overweight and Obese Adolescents: A Systematic Review and Meta-Analysis of Controlled Studies. *Int J Environ Res Public Health*. 2020;17(8).
50. Koskela-Staples NC, Yourell JL, Fedele DA, Doty J. Physical Activity Engagement: Perspectives From Adolescents With Comorbid Asthma and Overweight/Obesity and Their Caregivers. *J Pediatr Psychol*. 2023;48(8):707-19.
51. Jerome GJ, Fink T, Brady T, Young DR, Dickerson FB, Goldsholl S, et al. Physical Activity Levels and Screen Time among Youth with Overweight/Obesity Using Mental Health Services. *Int J Environ Res Public Health*. 2022;19(4).
52. Grossman DC, Bibbins-Domingo K, Curry SJ, Barry MJ, Davidson KW, Doubeni CA, et al. Screening for Obesity in Children and Adolescents: US Preventive Services Task Force Recommendation Statement. *JAMA*. 2017;317(23):2417-26.
53. Summary of the clinical practice guideline for multicomponent behavioral treatment of obesity and overweight in children and adolescents. *Am Psychol*. 2020;75(2):178-88.
54. Jones HM, Al-Khudairy L, Melendez-Torres GJ, Oyebo O. Viewpoints of adolescents with overweight and obesity attending lifestyle obesity treatment interventions: a qualitative systematic review. *Obes Rev*. 2019;20(1):156-69.

55. Ojeda-Rodríguez A, Zazpe I, Morell-Azanza L, Chueca MJ, Azcona-Sanjulian MC, Marti A. Improved Diet Quality and Nutrient Adequacy in Children and Adolescents with Abdominal Obesity after a Lifestyle Intervention. *Nutrients*. 2018;10(10).
56. Elvsaa I, Giske L, Fure B, Juvet LK. Multicomponent Lifestyle Interventions for Treating Overweight and Obesity in Children and Adolescents: A Systematic Review and Meta-Analyses. *J Obes*. 2017;2017:5021902.
57. O'Connor EA, Evans CV, Burda BU, Walsh ES, Eder M, Lozano P. Screening for Obesity and Intervention for Weight Management in Children and Adolescents: Evidence Report and Systematic Review for the US Preventive Services Task Force. *JAMA*. 2017;317(23):2427-44.
58. Srivastav P, K V, Bhat HV, Broadbent S. Effect of a pragmatic lifestyle modification intervention on physical activity levels and body mass index among obese and overweight adolescents in Udupi, India: a cluster randomized trial. *F1000Res*. 2024;13:859.
59. Srivastav P, K V, Bhat VH, Broadbent S. Structured, multifactorial randomised controlled intervention to investigate physical activity levels, body composition and diet in obese and overweight adolescents. *BMJ Open*. 2021;11(3):e044895.
60. Amiri P, Jalali-Farahani S, Zarkesh M, Gharibzadeh S, Hedayati M, Azizi F. Behavioral Interventions for Weight Management in Overweight and Obese Adolescents: A Comparison Between a Motivation-based Educational Program and Conventional Dietary Counseling. *Int J Endocrinol Metab*. 2020;18(1):e88192.
61. van der Kruk JJ, Kortekaas F, Lucas C, Jager-Wittenaar H. Obesity: a systematic review on parental involvement in long-term European childhood weight control interventions with a nutritional focus. *Obes Rev*. 2013;14(9):745-60.
62. Van Ryzin MJ, Nowicka P. Direct and indirect effects of a family-based intervention in early adolescence on parent-youth relationship quality, late adolescent health, and early adult obesity. *J Fam Psychol*. 2013;27(1):106-16.
63. Weihrauch-Blüher S, Kromeyer-Hauschild K, Graf C, Widhalm K, Korsten-Reck U, Jödicke B, et al. Current Guidelines for Obesity Prevention in Childhood and Adolescence. *Obes Facts*. 2018;11(3):263-76.
64. Liu Y, Ruan B, Jiang H, Le S, Liu Y, Ao X, et al. The Weight-loss Effect of GLP-1RAs Glucagon-Like Peptide-1 Receptor Agonists in Non-diabetic Individuals with Overweight or Obesity: A Systematic Review with Meta-Analysis and Trial Sequential Analysis of Randomized Controlled Trials. *Am J Clin Nutr*. 2023;118(3):614-26.
65. Wilding JPH, Batterham RL, Calanna S, Davies M, Van Gaal LF, Lingvay I, et al. Once-Weekly Semaglutide in Adults with Overweight or Obesity. *N Engl J Med*. 2021;384(11):989-1002.
66. Kadowaki T, Isendahl J, Khalid U, Lee SY, Nishida T, Ogawa W, et al. Semaglutide once a week in adults with overweight or obesity, with or without type 2 diabetes in an east Asian population (STEP 6): a randomised, double-blind, double-dummy, placebo-controlled, phase 3a trial. *Lancet Diabetes Endocrinol*. 2022;10(3):193-206.

67. Kelly AS, Auerbach P, Barrientos-Perez M, Gies I, Hale PM, Marcus C, et al. A Randomized, Controlled Trial of Liraglutide for Adolescents with Obesity. *N Engl J Med*. 2020;382(22):2117-28.
68. Bensignor MO, Bramante CT, Bomberg EM, Fox CK, Hale PM, Kelly AS, et al. Evaluating potential predictors of weight loss response to liraglutide in adolescents with obesity: A post hoc analysis of the randomized, placebo-controlled SCALE Teens trial. *Pediatr Obes*. 2023;18(9):e13061.
69. Zahmatkesh A, Sohoul M, Shojai S, Rohani P. The effect of orlistat in the treatment of non-alcoholic fatty liver in adolescents with overweight and obese. *Eur J Pediatr*. 2024;183(3):1173-82.
70. Lim F, Bellows BK, Tan SX, Aziz Z, Woo Baidal JA, Kelly AS, et al. Cost-Effectiveness of Pharmacotherapy for the Treatment of Obesity in Adolescents. *JAMA Netw Open*. 2023;6(8):e2329178.
71. Kelly AS. Current and future pharmacotherapies for obesity in children and adolescents. *Nat Rev Endocrinol*. 2023;19(9):534-41.
72. Pratt JSA, Browne A, Browne NT, Bruzoni M, Cohen M, Desai A, et al. ASMBS pediatric metabolic and bariatric surgery guidelines, 2018. *Surg Obes Relat Dis*. 2018;14(7):882-901.
73. O'Kane M, Parretti HM, Pinkney J, Welbourn R, Hughes CA, Mok J, et al. British Obesity and Metabolic Surgery Society Guidelines on perioperative and postoperative biochemical monitoring and micronutrient replacement for patients undergoing bariatric surgery-2020 update. *Obes Rev*. 2020;21(11):e13087.

Table 1. Summary of risk factors and possible pathogenesis of overweight and obesity in adolescents

Risk factors and pathogenesis	Details
Risk factors	Genetic: Polygenic variants interact with environmental factors. Lifestyle & Behavioural: Reduced physical activity, sedentary lifestyle, poor sleep, fast food, carbonated drinks, no daily vegetable intake, et al. Social & Environmental: Chronic psychosocial stress, food marketing to adolescents, shift to ultra-processed foods, loss of green spaces, increased motorized transport, tech changes, environmental contaminants.
Pathogenesis model	Complex multi-component model replacing the simple “calories in - calories out” model. Includes genetics, social-cultural backgrounds, obesogenic environment, and treatment barriers.
Regulatory systems of body weight	Energy Homeostasis: Unconscious system regulated by hypothalamus and brainstem. Receives long-term signals from adipose tissue (e.g., leptin) and short-term hunger/satiety signals from the gut. Cognitive-Emotional Control: Conscious system regulated by higher brain centres. Integrates homeostatic signals in the thalamus with environmental, experience, and emotion stimuli to control food intake.

Table 2. Summary of interventions for adolescents with overweight and obesity

Intervention type	Intervention strategies
Lifestyle interventions	Multicomponent lifestyle interventions are used as first-line treatment for anti-obesity interventions.
Dietary management	Energy restriction diet is beneficial for adolescents to control body weight, and reduce energy supply on the premise of ensuring the energy required for normal growth and development. Structured meal plans, portion control, and meal replacements can be used as possible weight management interventions. Dietary management emphasizes a balanced and healthy diet, and requires assessing and addressing modifiable risk factors in eating habits.
Physical activity	It is recommended 30 minutes moderate-intensity aerobic exercise at least 5 days a week, combined with strength or resistance training.
Behavioural management	Intensive healthy behaviour and lifestyle treatment requires patient and family collaboration with multidisciplinary treatment teams to conduct health education and skill training. Motivational home-based or school-based behavioural interventions are recommended.
Anti-obesity medications	Intensive health behaviour lifestyle treatment in combination with anti-obesity medications is recommended at the beginning of anti-obesity treatment in adolescents. Various anti-obesity medications, including glucagon-like peptide-1 receptor agonists, can reduce body weight and body mass index, but serious adverse events should be alert.
Metabolic surgery	Metabolic bariatric surgery must strictly adhere to clinical guideline indications and requires long-term multidisciplinary follow-up care.




Figure 1. Graphical abstract