

Supplementary Materials

Association between dietary fat intake and fatty acid profiles and hyperuricemia among Chinese adults: Results from the China Health and Nutrition Survey

Pengfeng Qu PhD¹, Yingying Jiao MM^{1,2}, Liusen Wang MM^{1,3}, Weiyi Li MPH^{1,3}, Hongru Jiang PhD^{1,3}, Jiguo Zhang MD^{1,3}, Huijun Wang MD^{1,3}, Bing Zhang MD^{1,3}, Junhua Han MD⁴, Aidong Liu MD^{1,3}, Zhihong Wang MD^{1,3}

¹Office of National Nutrition Plan, National Institute for Nutrition and Health, Chinese Center for Disease Control and Prevention, Beijing, China

²Hancheng Center for Disease Control and Prevention, Shaanxi, China

³Key Laboratory of Trace Elements and Nutrition, National Health Commission, Beijing, China

⁴Chinese Nutrition Society, Beijing, China

Corresponding Author:

Dr Zhihong Wang, National Institute for Nutrition and Health, Chinese Center for Disease Control and Prevention, 27 Nanwei Road, Beijing 100050, China.

Tel: +86-10-6623-7008.

Email: wangzh@ninh.chinacdc.cn


Dr Aidong Liu, National Institute for Nutrition and Health, Chinese Center for Disease Control and Prevention, 27 Nanwei Road, Beijing 100050, China.

Tel: +86-10-6623-7006.

Email: liuad@ninh.chinacdc.cn

Manuscript received 10 October 2025. Initial review completed 03 November 2025. Revision accepted 04 December 2025.

doi: 10.6133/apjcn.202604_35(2).0007



Supplementary Figure 1. Flow chart of study population selection