## Body mass index: is it an appropriate measure of obesity in Polynesians?

B Swinburn, MD, ChB, FRACP, P Craig, BSc, MHPEd, B Strauss, MBBS, PhD, FRACP, and R Daniel, DSM, MPH

Department of Community Health, University of Auckland, New Zealand; Royal Prince Alfred Hospital, Sydney; Monash University, Melbourne; Department of Health, Cook Islands.

The body mass index (BMI) is commonly used as an estimate of obesity with 20-25 kg/m<sup>2</sup> being considered normal. Polynesians, as individuals and populations, are often described as overweight or obese based on BMI criteria. We derived body fat measurements by bioimpedance methods in 129 adult Cook Island Polynesians and a representative sample of 505 adult Caucasian Australians using the same equation (Lukaski). As a group, the Cook Islanders were heavier (91.5  $\pm$  2.5 kg for men,  $78.9 \pm 2$  kg for women) compared to the Caucasians (80.7  $\pm$  1 kg for men, 66.4  $\pm$  1 kg for women; p < 0.0001 for both). Heights were similar and therefore the BMI values (kg/m<sup>2</sup>) were also higher in the Cook Islanders (29.6  $\pm$  0.7 and 29.8  $\pm$  0.7 versus 26.4  $\pm$  0.2 and  $25.3 \pm 0.2$ ; p<0.0001). Body fat, however, was lower in Cook Island men  $(22.0 \pm 1.0\% \text{ versus } 25.2 \pm 0.4\%;$ p<0.01) and similar in Cook Island women (31.2  $\pm$  1.0% versus  $33.0 \pm 0.3\%$ ). The following table of body fat estimations at equivalent BMI values shows that across the range Polynesians are leaner on a weight for weight basis:

	Body fat (%)		Body fat (%)	
	for males		for females	
At BMI	Polynesian	Caucasian	Polynesian	Caucasian
$20 \text{ kg/m}^2$	4.9	7.1	15.9	25.9
$25 \text{ kg/m}^2$	13.7	21.4	23.7	32.5
$30 \text{ kg/m}^2$	22.5	35.7	31.6	39.2
$40 \text{ kg/m}^2$	40	64.2	47.2	52.5

In conclusion, at any given BMI Polynesian men and women are leaner than Caucasians. If these data are confirmed with more definitive body composition studies, the BMI definitions may need to be altered for Polynesians, in which case a BMI of up to about 30 kg/m² could be considered normal.